

Fertilizer Research and Education Program

Final Report

A. Project Information

Project Title: Techniques to minimize nitrate loss from the root zone during managed aquifer recharge

Project leaders: Anthony Toby O'Geen and Helen Dahlke

Grant Number: 20-1019-000-SA

Project Duration: Start Date: 01/01/2021, End Date: 12/31/2025

Email and Phone (Lead PI): atogeen@ucdavis.edu, 530-752-2155

Report Type: Final Report

B. Abstract

Agricultural management of floodwater (Flood-MAR) is a practice where floodwaters are applied to agricultural fields to recharge groundwater. Flood-MAR may harm groundwater quality by leaching soil nitrate (NO_3^-) into groundwater. This modeling study evaluated Flood-MAR NO_3^- leaching risk in different climates and soil textures in California through multi-decadal simulations using the Root Zone Water Quality Model (RZWQM2). It evaluated whether Flood-MAR timing strategies (early vs late season irrigation application strategies) influenced risk. Flood-MAR had near negligible NO_3^- leaching risk in locations with rainfall $> 400 \text{ mm yr}^{-1}$. At drier locations, Flood-MAR NO_3^- leaching risk was highest (especially in loamy soils) because low rainfall allowed NO_3^- to build up in soils from year to year. Different Flood-MAR timing strategies (early season vs late season), combined with variable pauses in water applications (3 vs. 7 vs. 21-day intervals) showed no difference in NO_3^- leaching. Infrequent Flood-MAR should be practiced with care in arid climates and especially after prolonged droughts coinciding with more limited irrigation water supplies that constrain salt-leaching practices all favoring residual NO_3^- accumulation.

C. Introduction

The agricultural management of floodwaters (Flood-MAR) is of broad interest as a tool to recharge aquifers, offset groundwater pumping, and mitigate downstream flood risk (CDWR, 2018). This practice involves flooding agricultural fields during the winter and spring, and in some cases early summer, when rivers can reach high magnitude flows during wet years as a result of atmospheric river events and snowmelt (Kocis and Dahlke, 2017). Novel approaches are necessary to sustain irrigated agriculture in the face of multiple challenges. These include new public policy constraints on groundwater use, such as the Sustainable Groundwater Management Act requiring groundwater sustainability plans in overdrafted basins (CDWR, 2022), and additional challenges imposed by climate change, such as more intense precipitation whiplash

(Swain et al., 2018) and decreased mountain snowpack, a natural reservoir for California's irrigated agriculture that will diminish with warming (Qin et al. 2020).

However, there are concerns to be analyzed before Flood-MAR can be safely implemented, such as contamination of groundwater by flushing out residual soil nitrate (NO_3^-) during crop dormancy or fallowing at times that coincide with high magnitude river flows. Our research assumed an overarching hypothesis that Flood-MAR would enhance NO_3^- leaching compared to no Flood-MAR (business-as-usual), following the concern that Flood-MAR would mobilize residual NO_3^- in fertilized agroecosystems (Levintal et al., 2022). For example, in relatively deep vadose zone samples under California citrus orchard fertilizer trials, Harter et al. (2005) reported residual nitrate levels of 218-477 $\text{kg NO}_3^- \text{-N ha}^{-1}$ in 16-m deep cores. Bachand et al. (2014) measured residual NO_3^- in western Fresno County fields (San Joaquin Valley, California) where Flood-MAR had been practiced for one season and observed a residual NO_3^- quantity less than 15%, on average, of the soil profile NO_3^- (0-3 m) in wine grape control fields. In a deep soil coring study (0-30 m) across 36 study sites (12 fields) in the same region, Waterhouse et al. (2020) found relatively high residual NO_3^- across tomatoes and almonds (737-966 and 420-1,240 $\text{kg NO}_3^- \text{-N ha}^{-1}$, respectively) with a lower range for wine grapes where some growers tend to apply less fertilizer (75-1,371 $\text{kg NO}_3^- \text{-N ha}^{-1}$), but overall still demonstrating the risk for mobilizing residual NO_3^- with Flood-MAR.

D. Objectives

The overall objective of this project was to identify best management strategies for Flood-MAR to reduce the risk of groundwater nitrate contamination. These strategies are informed by trends in nitrogen cycling during the dormant season and the effects of flooding agricultural fields. These strategies will consider duration of water application, frequency of water applications and the associated periods of anaerobic and aerobic conditions which drive nitrogen transformations. was to simulate Flood-

The specific research objective was to evaluate scenarios of contrasting seasonal timing and frequency of Flood-MAR as strategies to minimize NO_3^- leaching risk, by leveraging the Root Zone Water Quality Model (RZWQM2), a widely validated tool developed and maintained by a team of USDA Agricultural Research Service scientists. We explored the primary controls (climate and soil properties) influencing the interaction of Flood-MAR with the N-cycle. RZWQM2 was used to simulate a multi-decadal business-as-usual scenario and five different Flood-MAR approaches in the context of a fertilized (250 $\text{kg N ha}^{-1} \text{ yr}^{-1}$) and irrigated maize agroecosystem with winter fallows across different representative soils and climates, as described below.

We explored the following research questions:

1. *Does Flood-MAR pose greater NO_3^- leaching risk when practiced early in the dormant season vs later in the season when conditions are warmer.*
2. How does the frequency of water application influence NO_3^- leaching risk?
3. How does soil texture influence NO_3^- leaching risk?

E. Methods

2.1 Overview of RZWQM2

RZWQM2 is a comprehensive process model that simulates fluxes of soil water and heat, organic matter cycling, and biogeochemical transport of N, linked to crop growth and nutrient and water uptake models, all paired to an agriculture management module (Ahuja et al., 2000). RZWQM2 simulations were performed on soil profiles (152 to 218-cm deep) with properties defined by horizon based on the Soil Survey Geographic database (SSURGO). RZWQM2 then automatically discretizes these user-defined soil horizons into multiple numerical nodes per horizon to simulate 1-D soil water and heat transport. Dynamic (transient) simulations spanned a 37-year simulation period with user-defined daily weather and management data, but variable time-steps depended on numerical solution convergence. Simulated initial conditions, (SIC), for RZWQM2 state parameters (water content, nitrate, SOM, etc.) were obtained by taking the end-of-simulation-time state of these parameters in an initial 37-year “spin-up” scenario.

2.1.1 Parameterization of RZWQM2

For this study, RZWQM2 modeling included 33 Central Valley soil series representing distinct taxonomic family particle-size classes (see section 2.3) and five different 37-year climate records obtained from the California Irrigation Management System (Table 1; see appendix), spanning a precipitation gradient from relatively wet-to-dry in space and time (n=165 unique soil-climate scenarios). The default RZWQM2 biogeochemical rate constants affecting N dynamics (e.g., nitrification, denitrification, etc.) were used across all scenarios. Initial values for SOM pool sizes, microbial populations, crop litter, and soil profile NO₃⁻ were allowed to vary by soil × climate combination and were established by using end-of-run values from the 37-year SIC run of each unique soil × climate modeling combination (33 soils × 5 climates =165).

In RZWQM2 simulations, Flood-MAR was practiced during the 10-wettest water years (Oct-Sep) of each specific 37-year climate record, applying 600-cm additional water (60 cm yr⁻¹) through Flood-MAR. During a Flood-MAR year, four 15-cm water applications were made in either January or March, using an application frequency of either 3- or 7-day intervals (Table 2). A fifth scenario tested applications from January-March with 21-day intervals. For fine textured soils with slower saturated hydraulic conductivities, Flood-MAR was practiced up to 9 continuous days to achieve 15-cm applied water in each event, depending on the soil’s depth-weighted hydraulic conductivity, using 3-, 7-, or 21-days between these continuous periods of Flood-MAR, depending on the scenario. RZWQM2 can mimic surface ponding by adjusting the Maximum surface storage parameter. In this study, in order to realistically represent FloodMAR, time periods to achieve 15-cm infiltrated water had to be determined *a priori* as shown in Table 2.

Table 2. Summary of conditions simulated using the Root Zone Water Quality Model.

Scenario	Flood-MAR application Frequency	Season
BAU Business as usual	NA	NA
High frequency-late	15 cm H ₂ O applied every 3 days	March
High Frequency-early	15 cm H ₂ O applied every 3 days	January
Low Frequency-late	15 cm H ₂ O applied every 7 days	March
Low Frequency-early	15 cm H ₂ O applied every 7 days	January
Extended Frequency	15 cm H ₂ O applied every 21 days	January through March

*Each Flood-MAR scenario consisted of 4 applications of 15 cm of water performed during the 10 wettest years of the 37-yr simulation period. BAU received no Flood-MAR treatment.

2.1.2 Simulated cropping system details

The objective for the project here was to have a stable crop growth model with reliable, consistent plant growth and N accumulation each year across all soil × climate scenarios, accepting that some minor differences would be expected. Ideally, similar yields and thus N removal through harvest would result in similar potentials for residual nitrate build-up due to less removal than applied across all simulated scenarios, to test the possible effects of Flood-MAR on additional leaching risk. For the model, planting occurred on April 15th every year, fertilization was applied as UAN in 4 splits with 50 kg total N applied in a preplant on April 15th, and the remainder divided evenly as 66.7 kg N applications on June 1st, July 1st, and July 21st. UAN applications were entered into RZWQM2 input files as appropriate proportions of NH₄, NO₃⁻, and urea. Irrigation was managed using the automated option in RZWQM2, where irrigation is triggered when plant available water is depleted to some allowable percentage threshold, called “allowable depletion,” and then the soil is refilled by irrigation back to field capacity. Allowable depletion was set at 60% during the first 20 days of crop growth, then reduced to 50% until harvest. This typically resulted in an irrigation frequency of several days during initial stand establishment and then every 5-14 days during the summer, depending on the soil’s water holding capacity. The simulated tillage depth was 15 cm with a tandem disk on April 15th before planting and then on September 15th and 16th after harvest with a bedder ridge operation on September 17th.

2.2 Climate data

Five stations from the California Irrigation Management Information System (CIMIS) were chosen on a wet-to-dry precipitation gradient (569 to 159 mm yr⁻¹) with at least a 37-year, largely intact record to provide climate data as input to RZWQM2 simulations (Table 1). Daily aggregated weather data were downloaded using the *cimir* package in R software (Koochafkan, 2021) and then subjected to in-house R software functions for visualizing anomalies and also automated quality-control checks, error correction, and gap-filling.

2.3 Soil data

33 soil profile datasets from SSURGO were used to parameterize the 990 RZWQM2 scenarios (1 business-as-usual and 5 Flood-MAR scenarios across 165 soil × climate conditions). The soil data objective was to represent widespread soils without restrictive horizons, spanning a broad textural gradient, and to generally test whether differences in inherent soil physical properties affected NO₃⁻ leaching response to Flood-MAR. The SSURGO sampling strategy was based initially on a soil classification study that defined seven generalized soil regions in California. Of these seven, three typically well-drained soil regions varying in texture were used as sampling targets, including coarse soils with no restrictions, loamy soils with no restrictions, and, a fine-textured endmember, shrink-swell soils (Devine et al. 2021).

2.4 Definition of “additional NO₃⁻ leaching risk”

For each unique Flood-MAR scenario (33 soils × 5 climates × 5 Flood-MAR timing strategies = 825 Flood-MAR scenarios), an “additional NO₃⁻ leaching risk” was calculated by subtraction of the total nitrate leached from its respective business-as-usual scenario that used the same soil and climate (BAU for 33 soils × 5 climate combinations = 165 BAU scenarios). This relative “additional NO₃⁻ leaching risk” term is used throughout the Results and Discussion (section 3) and acknowledges that the study is not attempting to predict any absolute amount of NO₃⁻ leaching for a given place or time. The purpose of the BAU scenario is to serve as a control to measure the extent to which Flood-MAR impacts NO₃⁻ leaching. In other words, interpretation of this study’s results are meant to ascertain relative risks of practicing Flood-MAR across different soils and climates in the Central Valley.

F. Data/Results and Discussion

3.1 Wetter climates prevent accumulation of residual NO₃⁻ in business-as-usual scenarios

Under BAU scenarios, the end-of-run residual soil NO₃⁻ largely stabilized across generalized taxonomic family particle-size classes in each climate modelled (Figure 1). This is inferred by the similarity in residual nitrate levels between the SIC (after 37 years of “spin-up”) and at the end of the 37-year BAU simulation (Figure 1). This suggests that, for each unique soil and climate combination, the SIC represent the appropriate conditions for the dynamic steady-state residual NO₃⁻ in the modeled agroecosystem, thus allowing for a robust test of the effect of Flood-MAR on additional NO₃⁻ leaching risk. Whole soil residual NO₃⁻ in wetter climates (Durham and Davis) were typically 60-100 kg N ha⁻¹ in the BAU scenario (Figure 1). Thus, climates with typical precipitation amounts of > 400 mm yr⁻¹ (Table 1) were sufficient to leach soils in this simulated, fertilized agroecosystem, suggesting that Flood-MAR practiced in wetter Central Valley climates is of relatively lower additional NO₃⁻ leaching risk (Figure 2). Modeled residual NO₃⁻ levels and the relationship with mean annual precipitation generally corresponded with levels found in the literature, although we were unable to find data for California. In Minnesota as an example, residual NO₃⁻-N under corn production ranged from around 100 kg N ha⁻¹ at locations with relatively low precipitation (400 mm yr) to 42 kg N ha⁻¹ at locations with high precipitation (891 mm yr) (Yadav, 1997). In addition, a 4-year trial in corn in eastern Colorado showed gradual build-up (23 kg/ha NO₃-N) of residual nitrate

in plots that were sub-optimally irrigated (46 cm of applied water) compared to half as much nitrate build-up (13 kg/ha NO₃-N) in plots that were over irrigated (85 cm of applied water) (Ludwick et al., 1976).




Figure 1. Preliminary (end of simulated initial conditions) and final residual soil nitrate (in the business-as-usual scenario, no Flood-MAR) by three generalized taxonomic family particle-size classes: coarse, loamy, and fine (n=11 soils per class). Each plot shows the different climates modeled from wettest (Durham, 537 mm yr⁻¹) to driest (Shafter, 143 mm yr⁻¹). Preliminary is the end of a 37-year preliminary run defining simulated initial conditions (SIC). Final is the end of a subsequent 37-year run, using the SIC end-of-run values for initialization. Note the difference in Y-axis scale in wet vs drier sites.

3.2 Accumulation of residual NO₃⁻ in dry climates drives additional NO₃⁻ leaching

Residual soil NO₃⁻ increased by an order of magnitude in business-as-usual coarse and loamy soils, but not in fine soils from wettest-to-driest climates (Figure 1). Compared to coarse soils, loamy soils tended to accumulate even more residual soil NO₃⁻ and displayed more variance from wet-to-dry. In the second driest climate (Five Points), BAU residual NO₃⁻ exceeded 500 kg NO₃⁻-N ha⁻¹ in 5 of 11 coarse soils (interquartile range: 277-686 kg NO₃⁻-N ha⁻¹); 8 of 11 loamy soils (interquartile range: 498-1,493 kg NO₃⁻-N ha⁻¹); and 2 of 11 fine soils (interquartile range: 77-301 kg NO₃⁻-N ha⁻¹). In the driest climate (Shafter), BAU residual NO₃⁻ exceeded 500 kg NO₃⁻-N ha⁻¹ in 7 of 11 coarse soils (interquartile range: 370-882 kg NO₃⁻-N ha⁻¹); 10 of 11 loamy soils (interquartile range: 586-1,860 kg NO₃⁻-N ha⁻¹); and 2 of 11 fine soils (interquartile range: 137-342 kg NO₃⁻-N ha⁻¹). In the driest climate, extreme values of residual NO₃⁻

accumulated in 2 of 11 coarse soils (1,693-2,249 kg NO₃-N ha⁻¹) and 4 of 11 loamy soils (1,831-3,145 kg NO₃-N ha⁻¹), but the most extreme fine soil accumulated far less (887 kg NO₃-N ha⁻¹).




Figure 2. Effect on 37-year, cumulative nitrate leached of different Flood-MAR strategies, including 3-day (d) and 7-d frequency events practiced either in January (Jan) or March (Mar) or 21-d frequency events spanning early Jan thru late Mar. Additional nitrate leached is relative to each soil's business-as-usual run (no Flood-MAR). Each plot is grouped by three generalized taxonomic family particle-size classes: coarse; loamy; and fine (n=11 soils per class); and shows the different climates modeled from wettest (Durham) to driest (Shafter). See Table 1 in appendix for descriptions of climates. Note the difference in Y-axis scale in wet vs drier sites.

Field studies in the region have demonstrated similar findings to this modeling exercise where soil profile nitrate was shown to accumulate during dry years and leached into the vadose zone during wet years (Liang et al., 1991; Rajj-Hoffman et al., 2024). Soil coring studies from the southern portions of the San Joaquin Valley, California, which correspond with the driest sites in this study, show residual NO₃⁻ masses similar to- but slightly lower (180 to 269 kg/ha) than the range in values derived from these RZWQM2 simulations of fertilized maize (Harter et al., 2005; Bachand et al., 2014; Waterhouse et

al., 2021). These studies sampled sites growing almonds, nectarine, tomato, grapes and wheat, and did not span the gradient in texture modeled here.

In drier climates, loamy soils tended to present the greatest possibility of risk of additional NO_3^- leaching with Flood-MAR (Figure 2). In the driest climate (Shafter), 4 of 11 loamy soils leached >3000 kg additional NO_3^- -N ha^{-1} over 37 years under the 21-day frequency Flood-MAR scenario, while the median flux was 1,270 kg additional NO_3^- -N ha^{-1} (interquartile range: 665-3,347 kg additional NO_3^- -N leached ha^{-1}). Although, coarse and fine textured soil groups only leached marginally less NO_3^- (median) compared to loamy soils during Flood-MAR treatments (Figure 2), the interquartile ranges of coarse and fine textural groups were much narrower: 695-1,007 and 622-1,214 kg additional NO_3^- -N leached ha^{-1} , respectively. In the second driest climate (Five Points), 3 of 11 loamy soils leached >3000 kg additional NO_3^- -N ha^{-1} with median fluxes of 990 kg additional NO_3^- -N ha^{-1} (interquartile range: 609-2,360 kg additional NO_3^- -N leached ha^{-1}). Similar to the driest climate, interquartile ranges of coarse and fine soil groups were much narrower: 499-764 and 532-1,103 kg additional NO_3^- -N leached ha^{-1} , respectively.

Although there are several facets of the N-cycle which can complicate this risk assessment (Figures 3 and 4; S1-S4), the most direct mechanistic explanation for additional NO_3^- leaching risk from loamy soils in the drier climates is due to their moderate level of microporosity and capacity to accumulate NO_3^- due to high water storage capacity. Loamy soils require more percolating water to leach effectively compared to coarse soils, explaining their conduciveness to residual NO_3^- accumulation in drier settings relative to coarse soils (Figure 1). These model findings are supported by deep core analysis from California field crops where a significant negative relationship was found between clay content in the soil profile (particle size control section) with NO_3^- concentration in the vadose zone (1.8-8 m; Lund et al., 1974). The study found low residual NO_3^- concentrations (< 5 $\mu\text{g/g}$) in the upper 1.8 m of sandy soils underlain by higher concentrations (20 $\mu\text{g/g}$) in the deep vadose zone. In contrast, comparatively high amounts of NO_3^- were found in the upper 1.8 m of loamy soils (peaking at 15 $\mu\text{g/g}$) with lower concentrations (5 $\mu\text{g/g}$) in the underlying deep vadose zone. Onsoy et al. (2005), in a 15 m deep, highly heterogeneous vadose zone, reported lowest NO_3^- -N concentrations in coarse-textured sediment facies relative to finer-texture sediments. Similarly, lab-based core leaching experiments have shown finer textured soils retained more NO_3^- -N compared to sandy soils (Gaines and Gaines, 1994).

While fine textured soils also have high water storage capacity, residual N build up was less (Figure 1) due to substantially higher denitrification (Figures 3 & 4). Fine textured soils have the potential to experience higher denitrification rates and/or longer episodes of denitrification due to longer durations of elevated water filled pore space (Groffman and Tiedje, 1988; Schindlbacher et al., 2004).

The RZWQM2 irrigation module uses an idealized irrigation schedule assuming perfect foresight by the farmer and uniform irrigation conditions: irrigation was triggered to apply water when a set threshold of plant available water was depleted. Upon triggering,

RZWQM2 applied a depth of water equal to refilling the soil's real-time, dynamic rooting depth to field capacity. This highly efficient irrigation decision approach had the effect of minimizing deep percolation, thus overestimating the accumulation of residual nitrate. To some extent, the higher NO_3^- mass leaching in the drier modelled climates would have been mitigated if this ideal business-as-usual irrigation scheme was less efficient or if winter water applications were used to mitigate salt build-up and thus lessened the impact of Flood-MAR on NO_3^- leaching concentrations.




Figure 3a-h. RZWQM2 simulated effects, comparing agricultural management of floodwaters (Flood-MAR) with business-as-usual (Control) on annualized N mass balance, hydrology, and water quality: a. Nitrate leached mass; b. Net mineralization; c. Harvested crop biomass N; d. Soil organic matter change affecting N balance; e. Volatilized N; f. Runoff; g. Deep percolation; and h. NO_3^- in deep percolation. Each plot is grouped by generalized taxonomic family particle-size classes: Coarse; Loamy; and Fine ($n=11$ per class). Crop system is $250 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ fertilized and irrigated maize in Parlier, CA (median precipitation= 278 mm yr^{-1}).

Under real, practical conditions, conventional irrigation management approaches recommend applying more water than necessary to avoid creating soil moisture stress. Furthermore, irrigation systems do not apply water uniformly and infiltration rates are inherently variable (Letey, 1985). For example, if 5 cm of applied water is needed to offset evapotranspiration, then a system with 80% irrigation distribution uniformity would need a 6.25 cm water application to prevent soil moisture stress. The consequence of this irrigation strategy is that most portions of a field will actually receive a depth of water greater than the amount necessary to refill the depleted soil moisture, resulting in deep percolation and NO_3^- leaching below the root zone as a consequence. A similar outcome is practiced in areas prone to salinity, where leaching fractions are used to manage salinity of root zones (Rhoades et al., 1973), now often post-harvest to avoid reducing nitrogen use efficiency. Applying more water to address salinity or irrigation distribution uniformity could complicate the interpretation of this study's results.

However, while such real-world practices may mean more deep percolation than this study suggests, it would also mean the Flood-MAR risk for additional NO_3^- leaching in this study is a conservatively high “book-end” estimate. Our results present a relatively conservative risk assessment and Flood-MAR would more clearly be a safe groundwater-quality mitigation tool in drier climates as well. Conversely, widespread use of subsurface drip irrigation throughout CA has been shown to limit root zone leaching and/or minimize leaching to concentrated areas surrounding the drip line (Hanson et al., 2008), leading to conditions more closely reflecting the simulated BAU here.




Figure 4. Effect on 37-year, cumulative denitrification of business-as-usual (no Flood-MAR) and various Flood-MAR strategies, including 3-day (d) and 7-d frequency events practiced either in January (Jan) or March (Mar) or 21-d frequency events spanning early Jan through late Mar. Each plot is grouped by three generalized taxonomic family particle-size classes: coarse; loamy; and fine (n=11 soils per class); and shows the different climates modeled from wettest (Durham) to driest (Shafter). See Table 1 in appendix for descriptions of climates.

3.3 Additional NO_3^- leaching risk in drier climates is mitigated by denitrification in finer textured soils and relative ease of leaching in coarse soils

The residual NO_3^- interquartile ranges of coarse and loamy texture groups tended to increasingly overlap from dry-to-driest (Figure 1). This convergence partly reflected the

denitrification capacity of loamy soils under high residual NO_3^- conditions (Figure 4), which contributed to mitigate additional NO_3^- leaching risk of loamy soils in drier climates (Figure 2). Although, loamy soils maintained an environment less favorable for denitrification compared to fine textured soils (D'Haene et al., 2003), as simulated by RZWQM2 (Figure 4).

In fine textured soils, denitrification limited the accumulation of residual soil NO_3^- to a greater extent, especially when dry climates favored build-up of NO_3^- due to less deep percolation (Figures 1, 3g, 4, S3g, S4g). Coarse soils displayed only negligible denitrification capacity, even under the driest climate (Figure 4). In fact, their capacity to accumulate some additional residual NO_3^- in the driest climate was reflected in the BAU run compared to the SIC run that assumed business-as-usual conditions (Figure 1). In addition to denitrification, the greater microporosity of fine soils physically limited business-as-usual NO_3^- leaching, especially in drier climates, making the notably lower total NO_3^- leached in Flood-MAR scenarios from fine soils more impactful on additional NO_3^- leaching risk due to Flood-MAR (Figure 3a, 5-7 appendix, S1a-S4a). This is in spite of the fact that deep percolation depths from fine soils were not markedly different compared to coarse and loamy soils in drier climates (Figure 3g, S3g-S4g). The retention of NO_3^- in soil under BAU, drier climate scenarios (Figures 5 and 7 see appendix) helps explain the similar relative responsiveness of coarse and fine soils to Flood-MAR in terms of additional NO_3^- leached (Figure 2). In the BAU scenario under the driest climate, coarse soils typically leached $1779 \text{ kg NO}_3\text{-N ha}^{-1}$ (interquartile range: $1305\text{-}1917 \text{ kg NO}_3\text{-N ha}^{-1}$), compared to $117 \text{ kg NO}_3\text{-N ha}^{-1}$ (interquartile range: $37\text{-}327 \text{ kg NO}_3\text{-N ha}^{-1}$) from fine soils (Figures 5 and 7, see appendix). Thus, this directly offset the Flood-MAR effect on leaching in coarse soils but exacerbated the NO_3^- leaching risk from fine soils, in spite of their high levels of denitrification and lower residual NO_3^- . Denitrification losses can be very high in fields with clay texture (Pratt et al., 1972), and the relationship between texture and denitrification has been documented where denitrification potential has been shown to be highest in fine textured soils, moderate in loamy soils, and low in coarse soils when water filled pore space exceeds 60% (D'Haene et al., 2003). A study of deep cores across a texture gradient in California found residual nitrate concentrations decreased with increasing clay content. The study suggests that while clay-rich soils have the potential to build-up in N via incomplete leaching, this N pool is removed by denitrification (Lund et al., 1974). Greater losses of NO_3^- via denitrification were simulated in fine (silt loam) compared to coarse (sandy loam) textured vadose zones during Flood-MAR using a reactive transport model (Waterhouse et al., 2021).

3.4 NO_3^- leaching risk “tipping point”

Comparing business-as-usual deep percolation rates versus additional NO_3^- risk highlights a clearly non-linear relationship to climate and, specifically, precipitation (Figure 8). The inflection or “tipping point” of this relationship is perhaps best exemplified by the Parlier climate (median annual precipitation: 278 mm yr^{-1}), where coarse soils showed less additional NO_3^- leaching risk compared to loamy and fine soils (Figure 2). The Parlier climate had ample precipitation to effectively leach most coarse soils in the initial business-as-usual scenario, preventing accumulation of residual NO_3^- ,

but not enough for most loamy soils (Figure 1). Accordingly, loamy soils leached more additional NO_3^- during Flood-MAR compared to coarse soils in the Parlier climate (Figure 2). Despite substantial denitrification limiting residual NO_3^- accumulation in fine soils in the Parlier climate (Figures 1 and 4), the typical fine textured soil leached slightly more additional NO_3^- (460 additional $\text{kg NO}_3\text{-N ha}^{-1}$) compared to loamy soils (395 additional $\text{kg NO}_3\text{-N ha}^{-1}$) (Figure 2). However, loamy soils arguably still had the greater additional NO_3^- leaching risk in the Parlier climate, due to their propensity for more extreme residual NO_3^- accumulation, despite the relatively high levels of business-as-usual NO_3^- leaching (3 of 11 soils developed $> 1,000 \text{ kg residual NO}_3\text{-N ha}^{-1}$ in Parlier, Figure 1). This was directly linked to additional NO_3^- leaching with Flood-MAR that was skewed towards more extreme fluxes in loamy soils (Figure 2). Close examination of RZWQM2 breakthrough curves for these more extreme loamy soils indicated that they did not generate substantial deep percolation until at least the second 15-cm Flood-MAR event. This explains the propensity of loamy soil to accumulate residual NO_3^- in business-as-usual scenarios in the just-dry-enough Parlier climate with not enough precipitation during even the wettest years to generate substantial deep percolation.




Figure 8. Relationship between total, 37-year deep percolation in the business-as-usual scenario (no Flood-MAR) and additional nitrate leached due to Flood-MAR practiced during the 10 wettest water years (Oct-Sep), applying $60 \text{ cm}^{-1} \text{ water yr}^{-1}$ across four applications on 21-day intervals. The vertical red line indicates the total Flood-MAR water applied over the 37-year period. Data shown across all climates modeled ($n=5$) and soils ($n=33$), symbolized by climate. Additional nitrate leached is relative to each soil's business-as-usual simulation in a particular climate. See Table 1 for description of climates.

3.5 Effect of Flood-MAR timing

Despite the marked increase in additional NO_3^- leaching with Flood-MAR in drier climates, no timing strategy was effective at appreciably mitigating this risk. However, there were consistent effects of late vs. early season Flood-MAR timing across different soils. Specifically, coarse and loamy soils experienced more additional NO_3^- leached during later season Flood-MAR scenarios (Figure 2, March 3-day and 7-day). Early season Flood-MAR experienced reduced additional NO_3^- leached because

mineralization rates are subdued at cooler temperatures (Miller and Geisseler, 2018). Moreover, Flood-MAR had the effect of reduced annual denitrification compared to BAU (Figure 4) by decreasing residual NO_3^- through leaching before the growing season when denitrification potential is greatest on account of warmer, more biogeochemically active soils. In addition to the amounts of O_2 and available C, residual NO_3^- concentration is a primary control of denitrification (Myrold and Tiedje, 1985).

3.6 Flood-MAR following multi-year droughts presents some risk

Across the 37-year simulation with Flood-MAR during the 10 wettest winters, median Flood-MAR scenarios tracked business-as-usual scenarios closely in wetter climates: the typical coarse soil leached just 43 and 45 kg additional NO_3^- -N ha^{-1} in Durham and Davis, respectively (Figure 5, see appendix); the typical loamy soil leached just 54 and 63 kg additional NO_3^- -N ha^{-1} in Durham and Davis (Figure 6, see appendix); and the typical fine soil leached 152 and 336 additional NO_3^- -N ha^{-1} in Durham and Davis, respectively (Figure 5).

However, in drier climates, residual NO_3^- accumulated during periodic droughts in the climate record (e.g. 2012-2016), which was then flushed as deep percolation during subsequent wet years (e.g. 2017) in the Flood-MAR scenarios on typical coarse and loamy soils (Figures 5 and 6, see appendix). Thus, the timing of Flood-MAR following multi-year droughts is of special concern in drier climates (median precipitation < 400 mm) and especially on more difficult-to-leach soils where high levels of residual NO_3^- accumulation are possible. It is during these dry years that residual NO_3^- would be accumulating, but relatively rare in these locations that wet year precipitation is sufficient to initiate substantial deep percolation capable of leaching soils effectively. If precipitation whiplash continues, strategies will be needed to deplete residual nitrate either via improved fertigation methods (higher nitrogen use efficiency), crop rotations or use of cover crops (Delgado et al., 1999).

G. Conclusions

Coarse soils are considered of greatest risk to NO_3^- leaching into groundwater when N-fertilizer applications exceed N-removal processes in an agroecosystem: export in crop biomass; accumulation in soil organic matter; volatilization; and denitrification (Figure 4). However, this truism did not hold up to evaluations of the effect of agricultural management of floodwater (Flood-MAR) on additional NO_3^- leaching risk. Simply put, except in the driest climates, precipitation is sufficient in California's Central Valley to leach residual NO_3^- , such that the additional NO_3^- leaching risk presented by Flood-MAR is typically lower in coarse soils compared to loamy soils.

Overall, this research suggests that Flood-MAR timing strategies, specifically seasonality and intervals between Flood-MAR events, are generally negligible in their effects on NO_3^- leaching risk in comparison to the risk of accumulating residual nitrate in dry climates. Nevertheless, results point to broader timing concerns for using Flood-MAR as a general strategy to recharge depleted aquifers. Most importantly, growers are advised to manage nitrogen carefully in dry years, monitor residual NO_3^- following successive dry years, especially in locations where natural deep percolation is already

typically limited by low rainfall. Residual NO_3^- may accumulate most rapidly during droughts, especially when growers face limited water resources that may typically result in lower-than-expected growing season soil flushing due to irrigation and leaching practices. However, under all scenarios, Flood-MAR is likely to noticeably reduce NO_3^- concentration in groundwater relative to business-as-usual over the long run.

H. Challenges

This study evaluated nitrate leaching risk of 33 different soil series with contrasting physical properties representative of commonly mapped soils across the region. The broad scale of the study domain, magnitude of time evaluated, and large number of Flood-MAR scenarios investigated made it impossible to calibrate and validate the model uniformly with field data. Thus, for the most part, model default conditions were used. This study only modeled one of the over 200 crops grown in California. Various crops will inevitably have different N requirements, irrigation strategies, rooting depths, and N use efficiencies, which will influence residual nitrate build-up. Although the residual N accumulation will differ among crops, this study is generalizable to any crop that has a dormant period over the winter months and some degree of residual nitrate accumulation. Despite these limitations, this study has value in presenting new testable research questions and management considerations such as the likelihood of denitrification during the dormant season, the potential build-up of residual nitrate in response to drought, effects of irrigation management and the potential impact of climate whiplash on residual NO_3^- , and the relevance of N-cycle process in the context of Flood-MAR. Findings are not intended to prescribe the ideal conditions for Flood-MAR, rather they justify the importance of documenting residual nitrate concentrations when considering this practice.

I. Project Impacts

Results shown here are relevant due to the potentially long-term implementation of various Flood-MAR regimes as climate resiliency and groundwater (supply) sustainability practices are expected to expand in the future (Marr et al., 2018). Simulations here pertain to losses of NO_3^- from the most dynamic and reactive portion of the unsaturated zone, the upper two meters of soil. Conservatively, assuming relatively low reactivity in the vadose zone below soil and furthermore assuming laterally widespread application of these simulated conditions, the NO_3^- concentration in leachate from soil would be similar when it becomes recharge at the water table. And, if that recharge has the same concentration across relatively large lateral extent, it would be expected that shallower (predominantly domestic) wells also experience concentrations of that magnitude, albeit after some period of delay corresponding to the travel time from the root zone to the water table and from the water table to the well (typically few to tens of years). In this section, we address two questions: what concentration of NO_3^- does the aforementioned additional losses of NO_3^- mass due to Flood-MAR represent? And, second, how is that concentration different from those under BAU conditions?

The difference in mass losses from soil range from few tens to several thousand kg N/ha over the simulated period (figs 5-7 see appendix). The main difference in water

percolation over that same period is approximately 600 cm across all scenarios (the amount of additional Flood-MAR applied). The drinking water limit of 10 mg NO₃⁻/L is equivalent to 1 kg N / (ha * 1 cm) or 10 kg N/ha in 10 cm of water (10 kg N in 0.1 ha-m of water) or 600 kg N/ha in 600 cm of water. From basic mass balance considerations, the *additional* water percolating from Flood-MAR does not exceed the nitrate maximum contaminant level (MCL), where the additional N leaching over the simulation period does not exceed 600 kg N/ha (Figure 8: almost all wet climate scenarios in Durham and Davis, some scenarios in intermediate climate in Parlier, few scenarios in the dry climate). Under dry climate conditions, where median N losses are on the order of twice 600 kg N/ha, the *additional* water percolating from Flood-MAR may be on the order of twice the MCL concentration, in some cases five times the MCL concentration (where the additional leaching is 3000 kg N/ha). Clearly, under the drier climate conditions, without adding additional practices such as improved nitrogen use efficiency, crop rotation, cover crops or higher Flood-MAR applications than 60 cm in the wettest winters, the percolation due to the additional Flood-MAR water cannot meet drinking water standards. In fact, modeled NO₃⁻ concentration in deep percolation waters receiving Flood-MAR treatment were reduced to equal the drinking water standard for most fine soil groups, while being slightly to moderately above the standard for loamy and coarse texture groups depending on location (Fig. 3h; S1-4).

The NO₃⁻ concentrations of deep percolation water in the BAU scenarios were higher in dry regions (Figures 3h). Moreover, NO₃⁻ concentrations of deep percolation water in the BAU scenarios can be estimated from the total mass of N leached under BAU conditions (no Flood-MAR; Figures 5 – 7) and the corresponding total BAU percolation (Figure 8). The total BAU NO₃⁻ leached was highest in coarse soils, ranging from 1000 kg N/ha under dry climates to over 1500 kg N/ha in wet climates (Figure 5). The corresponding percolation rates over the simulation period range from less than 20 to 250 cm in drier climates and from 300 to over 1200 cm in wet climates (Figure 8). Hence, under BAU conditions of drier climate, nitrate-N concentrations in (very slowly) percolating water exceed the MCL by nearly one to nearly two orders of magnitude (1500/250 to 1500/20 kg N/(ha*cm water)), while, under BAU conditions of wetter climates, nitrate-N concentrations in (rapidly) percolating water range from at the MCL to 5 five times above the MCL (1500/300 to 1500/1200 kg N/(ha * cm water)) under the simulated conditions.

Hence, under all climate conditions, the NO₃⁻ concentration in the *additional* percolating water due to Flood-MAR is significantly lower than the NO₃⁻ concentration in percolating water from BAU conditions. Due to the mixing of these two waters in the root zone and further in the deeper unsaturated zone, shallow groundwater, and along the vertical well screen of an extraction well, the net long-term effect of Flood-MAR is a significant reduction of NO₃⁻ concentration in shallow groundwater and, hence, in domestic and shallow public water supply water under dominant influence of agricultural return water recharge. Despite the fact that Flood-MAR leads to additional nitrate *mass* leaching, especially under dry climate conditions, the amount of Flood-MAR in the simulated scenarios is sufficient to dilute this additional nitrate mass to less than or slightly above the NO₃⁻ MCL depending on soil type and climate (Figure 3h). Pratt and others (1972)

arrived at a similar finding showing low [NO₃⁻] (below MCL) in deep leachate from orange orchards in the Central Valley due to the application of excess irrigation water to maintain low salt levels in soils.

The above considerations, while hypothetical, are highly relevant for consideration of the long-term impacts of Flood-MAR on well water NO₃⁻ concentrations, but will be further modified due to the fact that stream and incidental canal recharge with typically negligible NO₃⁻ concentrations play an important role particularly in the drier climate conditions of the southern Central Valley, leading to further dilution of irrigated lands return flow recharge NO₃⁻ in some wells. We note that, under the typical percolation and recharge rates of the drier climate scenarios (50 cm / 37 years) when compared to the wet climate scenarios (500 cm / 37 years, Figure 8), the source area of a well, at any given pumping rate, would be an order of magnitude larger in dry climate locations than in wet climate location, assuming the entire landscape was subject to the simulated scenario conditions.

J. Outreach Activities Summary

Event Name (1)	UC ANR Statewide meeting		
Presentation title	Decision support tools for soil and water resource management		
Location and date	Davis, CA 4/25/23		
Attendee demographics	Researchers, Extension experts		
	no	Number of participants	45

Event Name (2)	UC Davis, NRCS and UCCE soil data needs meeting		
Presentation title	Technical soil survey considerations to implement a new nitrate leaching hazard rating for the Groundwater nitrate leaching hazard index		
Location and date	Fresno CA		
Attendee demographics	NRCS staff, Cooperative Extension academics		
CCA/Grower Continuing Education Units (CEUs)	no	Number of participants	14

Event Name (3)	Southwest chapter Soil and Water Conservation Society		
Presentation title	Interactive Tools Supporting Soil and Water Resource Management, Lodi CA NRCS, state agency staff,		

	academics. Southwest chapter Soil and Water Conservation Society		
Location and date	Lodi, CA 11/1/23		
Attendee demographics	NRCS staff, UCCE academics, Consultants, growers CCA, PCAs		
CCA/Grower Continuing Education Units (CEUs)	yes	Number of participants	55

Event Name (4)	6th Western Groundwater Congress		
Presentation title	Estimating the impact of vadose zone heterogeneity on agricultural managed aquifer recharge: A combined experimental and modeling study		
Location and date	Burbank, CA 9/14/23		
Attendee demographics	Academics, consultants, agency staff		
CCA/Grower Continuing Education Units (CEUs)	no	Number of participants	45

Event Name (5)	California Extreme Precipitation Symposium		
Presentation title	Managed Aquifer Recharge as a Tool to Enhance Sustainable Groundwater Management in California.		
Location and date	Davis CA 6/27/23		
Attendee demographics	Academics, consultants, agency staff		
CCA/Grower Continuing Education Units (CEUs)	no	Number of participants	300

Event Name (6)	UC ANR statewide Conference		
Presentation title	Storing More Water Through Intentional Groundwater Recharge		
Location and date	Fresno, CA 4/25/23		
Attendee demographics	UC Cooperative extension academics, policy makers		
CCA/Grower Continuing Education Units (CEUs)	no	Number of participants	88

Event Name (7)	Madera County Almond Day		
----------------	--------------------------	--	--

Presentation title	Agricultural managed aquifer recharge: Plant response, hydrologic, and geochemical processes.		
Location and date	Madera, CA 3/23/23		
Attendee demographics	CCAs, PCAs, growers, consultants researchers		
CCA/Grower Continuing Education Units (CEUs)	yes	Number of participants	60

Event Name (8)	Dixon and Solano RCDs Groundwater workshop		
Presentation title	Nitrogen fate during agricultural managed aquifer recharge: Linking plant response, hydrologic, and geochemical processes		
Location and date	Dixon, CA 1/24/23		
Attendee demographics	Growers, Resource Conservation District staff, academics		
CCA/Grower Continuing Education Units (CEUs)	no	Number of participants	50

Event Name (9)	California Dairy Research Foundation		
Presentation title	AgMAR on working lands managed by dairies		
Location and date	Davis		
Attendee demographics	Key dairy stakeholders, academic advisors		
CCA/Grower Continuing Education Units (CEUs)	no	Number of participants	10

Event Name (10)	California Water & Environmental Modeling Forum		
Presentation title	Numerical Modeling to Assess Benefits of Efficient Nitrate Management and AgMAR on Groundwater		
Location and date	Folsom, 4/18/23		
Attendee demographics	Consultants, students, agency personnel		
CCA/Grower Continuing Education Units (CEUs)	no	Number of participants	60

Event Name (11)	Water Education Foundation		
Presentation title	Managed Aquifer Recharge		

Location and date	Central Valley Tour, 4/26/23		
Attendee demographics	Consultants, engineers, local, state, and federal agency personnel, irrigation and water district personnel, attorneys, environmental stakeholders		
CCA/Grower Continuing Education Units (CEUs)	no	Number of participants	50

Event Name (12)	Contemporary Groundwater Issues Council Annual Meeting		
Presentation title	Future of MAR, Water Management, and Water Infrastructure with Climate Uncertainty		
Location and date	Davis, CA 10/27/23		
Attendee demographics	California groundwater industry, state and regional decision-makers, leading groundwater academics in California		
CCA/Grower Continuing Education Units (CEUs)	no	Number of participants	49

Event Name (13)	DWR's Water Quality Monitoring Technical Advisory Committee		
Presentation title	Water quality in AgMAR		
Location and date	Online, 9/14/23		
Attendee demographics	Personnel from regulatory agencies, NGOs, academia, consultancies		
CCA/Grower Continuing Education Units (CEUs)	no	Number of participants	8

Event Name (14)	FREP/WPHA Conference		
Presentation title	Modeling Techniques to Minimize Nitrate Loss from the Root Zone During Managed Aquifer Recharge		
Location and date	Seaside, CA Oct. 30 th 2024		
Attendee demographics	CCA, PCAs, growers, consultants, researchers, CDFA staff, NRCS staff		
CCA/Grower Continuing Education Units (CEUs)	Nutrient Management: 5 Soil and Water Management: 3	Number of participants	~75

	Crop Management: 1 Professional Development: 0.5		
--	---	--	--

Event Name (15)	Updating USDA-NRCS on research from the California Soil Resource Lab		
Presentation title	Regional Modeling of Techniques to Limit Nitrate Leaching Risk During Flood-MAR		
Location and date	Davis CA Aug. 20th 2024		
Attendee demographics	USDA-NRCS area soil scientists		
CCA/Grower Continuing Education Units (CEUs)		Number of participants	10

K. References

Ahuja, L.R., Rojas, K.W., Hanson, J.D., Shaffer, M.J., Ma, L. (Eds.), 2000. Root Zone Water Quality Model. Water Resources Publications, Highland Ranch, CO.

Bachand, P.A.M., S. B. Roy, J. Choperena, D. Cameron, and W. R. Horwath, 2014. Implications of using on-farm flood flow capture to recharge groundwater and mitigate flood risks along the Kings River, CA. Environ. Sci. Technol. 48: 13601-13609. [dx.doi.org/10.1021/es501115c](https://doi.org/10.1021/es501115c) |

Bateman, E.J., E.M. Baggs, 2005. Contributions of nitrification and denitrification to N₂O emissions from soils at different water-filled pore space. Bio. Fert. Soil 41:379-388. <https://doi.org/10.1007/s00374-005-0858-3>

Cameira, M.R., Fernando, R.M., Ahuja, L.R., Ma, L., 2007. Using RZWQM to simulate the fate of nitrogen in field soil–crop environment in the Mediterranean region. Agr. Water Manage 90 (1-2), 121–136. <https://doi.org/10.1016/j.agwat.2007.03.002>.

CDWR (California Department of Water Resources), 2018. Flood-MAR: Using Flood Water for Managed Aquifer Recharge to Support Sustainable Water Resources. CDWR, Sacramento. https://cawaterlibrary.net/wp-content/uploads/2018/07/DWR_FloodMAR-White-Paper_06_2018_updated.pdf

CDWR (California Department of Water Resources), 2022. Groundwater Sustainability Plans. <https://water.ca.gov/Programs/Groundwater-Management/SGMA-Groundwater-Management/Groundwater-Sustainability-Plans>

Delgado, J.A., R.T. Sparks, R.F. Follett, J.L. Sharkoff, and R.R. Riggerbach. 1999. Use of winter cover crops to conserve water and water quality in the San Luis Valley of south

central Colorado. p. 125–142. *In* R. Lal (ed.) Soil quality and soil erosion. CRC Press, Boca Raton, FL.

Devine, S.M., K.L. Steenwerth and A. T. O’Geen. 2021. A Regional Soil Classification Framework to Improve Soil Health Diagnosis and Management. *Soil Science Society of America Journal*. 85: 361-378 <https://doi.org/10.1002/saj2.20200>

D’Haene, K., E. Moreels, S. DeNeve, B. Daguilar, P. Boeckx, G. Hofman and O. Van Cleemput, 2003. Soil properties influencing the denitrification potential of Flemish agricultural soils. *Biol. & Fert. Soils*. 38: 358-366.

Domagalski, J.L., Phillips, S.P., Bayless, E.R., Zamora, C., Kendall, C., Wildman, R.A., Hering, J.G., 2008. Influences of the unsaturated, saturated, and riparian zones on the transport of nitrate near the Merced River, California, USA. *Hydrogeol. J.* 16, 675–690. <https://doi.org/10.1007/s10040-007-0266-x>

Fang, Q. X., T. R. Green, L. Ma, R. W. Malone, R. H. Erskine, and L. R. Ahuja. 2010. Optimizing soil hydraulic parameters in RZWQM2 using automated calibration methods. *Soil Sci. Soc. Am. J.* 74: 1897-1913. <https://doi.org/10.2136/sssaj2009.0380>

FAO (Food and Agriculture Organization of the United Nations). 2021. World Food and Agriculture - Statistical Yearbook 2021. Rome. <https://doi.org/10.4060/cb4477en>

Gaines, T.P. and S.T. Gaines, 1994. Soil texture effect on nitrate leaching in soil percolates. *Comm. Soil Sci. Plant Anal.* 25:2561-2570 <https://doi.org/10.1080/00103629409369207>

Geisseler, D., Lazicki, P.A., Pettygrove, G.S., Ludwig, B., Bachand, P.A.M., Horwath, W.R., 2012. Nitrogen Dynamics in Irrigated Forage Systems Fertilized with Liquid Dairy Manure. *Agron. J.* 104(4), 897–907. <https://doi.org/10.2134/agronj2011.0362>

Green, W. H., & Ampt, G. A. 1911. Studies on soil physics: The flow of air and water through soils. *Journal of Agricultural Science*, 4(1), 1-24.

Groffman, P.M. and J.M. Tiedje, 1988. Denitrification hysteresis during wetting and drying cycles in soil. *Soil Sci. Soc. Am. J.* 52:1626-1629.

Hanson, B., J. Hopmans, and J. Simunek, 2008. Leaching with subsurface drip irrigation under saline, shallow groundwater conditions. *Vadose Zone J.* 7:810-818.

Harter, T., Onsoy, Y., Heeren, K., Denton, M., Weissmann, G., Hopmans, J., Horwath, W., 2005. Deep vadose zone hydrology demonstrates fate of nitrate in eastern San Joaquin Valley. *Calif. Agric.* 59, 124–132. <https://doi.org/10.3733/ca.v059n02p124>

Herbert, C. and Doll, P. 2019. Global assessment of current and future groundwater stress with a focus on transboundary aquifers. *Water Resources. Research*, 55, 4760–4784.

<https://doi.org/10.1029/2018WR023321>

Hoogenboom, G., C.H. Porter, K.J. Boote, V. Shelia, P.W. Wilkens, U. Singh, J.W. White, S. Asseng, J.I. Lizaso, L.P. Moreno, W. Pavan, R. Ogoshi, L.A. Hunt, G.Y. Tsuji, and J.W. Jones. 2019. The DSSAT crop modeling ecosystem. In: p.173-216 [K.J. Boote, editor] *Advances in Crop Modeling for a Sustainable Agriculture*. Burleigh Dodds Science Publishing, Cambridge, United Kingdom
(<https://dx.doi.org/10.19103/AS.2019.0061.10>).

Kisekka, I., Schlegel, A., Ma, L., Gowda, P., Vara Prasad, P. 2017. Optimizing preplant irrigation for maize under limited water in the high plains. *Agricultural Water Management*. 187:154-163.

Kocis, T.N., Dahlke, H.E., 2017. Availability of high-magnitude streamflow for groundwater banking in the Central Valley, California. *Environ. Res. Lett.* 12(8), 084009.
<https://doi.org/10.1088/1748-9326/aa7b1b>

Koohafkan, M., 2021. cimir: Interface to the CIMIS Web API. R package version 0.4-1.
<https://CRAN.R-project.org/package=cimir>

Letey J. 1985. Irrigation uniformity as related to optimum crop production-Additional research is needed. *Irrig. Sci.* 6 253-263.
<https://link.springer.com/article/10.1007/BF00262470>

Levintal, E., Kniffin, M.L., Ganot, Y., Marwaha, N., Murphy, N.P., Dahlke, H.E., 2022. Agricultural managed aquifer recharge (Ag-MAR)—a method for sustainable groundwater management: A review. *Crit. Rev. Env. Sci. Technol.*
<https://doi.org/10.1080/10643389.2022.2050160>

Liang, B.C., M. Remillard, A.F. MacKenzie, 1991. Influence of fertilizer, irrigation, and non-growing season precipitation on soil nitrate-nitrogen under corn. *J. Environ. Qual.* 20: 123-128. <https://doi.org/10.2134/jeq1991.00472425002000010019x>

Liu, P.W., Famiglietti, J.S., Purdy, A.J. Adams, K.H., McEvoy, A.L., Reager, J.T., Bindlish, R. Wiese, D.N., David, C.H., Rodell, M., 2022. Groundwater depletion in California's Central Valley accelerates during megadrought. *Nat. Commun.* 13, 7825.
<https://doi.org/10.1038/s41467-022-35582-x>

Lund, L.J., D.C. Adriano, and P.F. Pratt. 1974. Nitrate concentrations in deep soil cores as related to soil profile characteristics. *J. Environ. Qual.* 3:78-82.
<https://doi.org/10.2134/jeq1974.00472425000300010021x>

- Ma, L., Shaffer, M.J., Boyd, J.K., Waskom, R., Ahuja, L.R., Rojas, K.W., Xu, C., 1998. Manure management in an irrigated silage corn field: experiment and modeling, *Soil Sci. Soc. Am. J.* 62, 1006–1017. <https://doi.org/10.2136/sssaj1998.03615995006200040023x>
- Ma, L., Hoogenboom, G., Ahuja, L.R., Ascough, J.C., Saseendran, S.A., 2006. Evaluation of the RZWQM-CERES-Maize hybrid model for maize production. *Agricultural Systems* 87(3), 274–295. <https://doi.org/10.1016/j.agsy.2005.02.001>
- Ma, L., Ahuja, L.R., Saseendran, S.A., Malone, R.W., Green, T.R., Nolan, B.T., Bartling, P.N.S., Flerchinger, G.N., Boote, K.J., Hoogenboom, G., 2011. A Protocol for Parameterization and Calibration of RZWQM2 in Field Research, in: Ahuja, L.R., Ma, L., (Eds.), *Methods of Introducing System Models into Agricultural Research*, ASA, CSSA, SSSA, Madison, WI, pp. 1–64. <https://doi.org/10.2134/advagricsystmodel2.c1>
- Marr, J., D. Arrate, R. Maendly, D. Dhillon, and S. Stygar, 2018. FLOOD-MAR: Using flood water for managed aquifer recharge to support sustainable water resources. California Department of Water Resources White paper. <https://water.ca.gov/Programs/All-Programs/Flood-MAR>
- Mekala, C. and I.M. Nambi, 2017. Understanding the hydrologic control of N cycle: Effect of water filled pore space on heterotrophic nitrification, denitrification and dissimilatory nitrate reduction to ammonium mechanisms in unsaturated soils. *J Contaminant Hydrol.* 202: 11-22. <https://doi.org/10.1016/j.jconhyd.2017.04.005>
- Miller, K.S. and D. Geisseler, 2018. Temperature sensitivity of nitrogen mineralization in agricultural soils. *Biol. & Fert. Soils* 54:853-860. <https://doi.org/10.1007/s00374-018-1309-2>
- Murphy, N.P., Waterhouse, H., and H.E. Dahlke, 2021. Influence of agricultural managed aquifer recharge on nitrate transport: The role of soil texture and flooding frequency. *Vadose Zone J.* 20(5), 1–16. <https://doi.org/10.1002/vzj2.20150>
- Myrold, D.D. and J. M. Tiedje, 1985. Diffusional constraints on denitrification in soil. *Soil Sci. Soc. Am. J.* 49: 651-657 <https://doi.org/10.2136/sssaj1985.03615995004900030025x>
- Nelson, R.L., 2012. Assessing local planning to control groundwater depletion: California as a microcosm of global issues, *Water Resour. Res.* 48, W01502. <https://doi.org/10.1029/2011WR010927>
- Nimah N. M., and R. J. Hanks, 1973. Model for Estimating Soil Water, Plant, and Atmospheric Interrelations, I. Description and Sensitivity, *Soil Science Society of America Proceedings*, 37: 522-527.

- Onsoy, Y.S., T. Harter, T.R. Ginn and W.R. Horwath, 2005. Spatial variability and transport of nitrate in a deep alluvial vadose zone. *Vadose Zone J.* 4:41-54.
- Or, D., B.F. Smets, J.M. Wraith, A. Dechesne and S.P. Friedman, 2007. Physical constraints affecting bacterial habitats and activity in unsaturated porous media – a review. *Advances in Water Resources* 30:1505-1527
<https://doi.org/10.1016/j.advwatres.2006.05.025>
- Pratt, P.F., Jones, W.W., Hunsaker, V.E., 1972. Nitrate in deep soil profiles in relation to fertilizer rates and leaching volume. *Environ. Qual.* 1, 97–102.
<https://doi.org/10.2134/jeq1972.00472425000100010024x>
- Qin, Y., Abatzoglou, J.T., Siebert, S., Huning, L.S., AghaKouchak, A., Mankin, J.S., Hong, C., Tong, D., Davis, S.J., Mueller, N.D., 2020. Agricultural risks from changing snowmelt. *Nat. Clim. Change* 10, 459–465. <https://doi.org/10.1038/s41558-020-0746-8>
- Raij-Hoffman, I., O. Dahan, H. E. Dahlke, T. Harter, I. Kisekka. 2024. Assessing Nitrate Leaching During Drought and Extreme Precipitation: Exploring Deep Vadose-Zone Monitoring, Groundwater Observations, and Field Mass Balance. *Water Resour. Res.* 60, e2024WR037973. <https://doi.org/10.1029/2024WR037973>
- Rawls, W.J., D.L. Brakensiek, and K.E. Saxton. 1982. Estimation of soil-water properties. *Trans. Am. Soc. Agric. Eng.* 25:1316–1320, 1328. doi: 10.13031/2013.33720
- Rhoades, J.D., R.D. Ingvalson, J.M. Tucker and M. Clark 1973. Salts in irrigation drainage waters: I. Effects of irrigation water composition leaching fraction, and time of year on the salt compositions of irrigation drainage waters *Soil, Sci. Soc. Am. Proc.* 37: 770-774.
- Richards, L.A. 1931. Capillary Conduction of Liquids through Porous Mediums. *Journal of Applied Physics*, 1: 318-333. <http://dx.doi.org/10.1063/1.1745010>
- Schindlbacher, A., S. Zechmeister-Boltenstern; K. Butterbach-Bahl, 2004. Effects of soil moisture and temperature on NO, N₂O, and N₂O emissions from European forest soils. *J. Geophysical Res.* 109D17.
- Siebert, S., Burke, J., Faures, J.M., Frenken, K., Hoogeveen, J., Döll, P., Portmann, F.T., 2010. Groundwater use for irrigation – a global inventory, *Hydrol. Earth Syst. Sci.* 14, 1863–1880. <https://doi.org/10.5194/hess-14-1863-2010>
- Soil Survey Staff, 1999. *Soil Taxonomy: A basic system of soil classification for making and interpreting soil surveys*. Second ed. USDA-NRCS Agricultural Handbook No. 436.
- Swain, D.L., Langenbrunner, B., Neelin J.D., Hall, A., 2018. Increasing precipitation volatility in twenty-first-century California *Nat. Clim. Change* 8, 427–33.
<https://doi.org/10.1038/s41558-020-0746-8>

Waterhouse, H., Bachand, S., Mountjoy, D., Choperena, J., Bachand, P., Dahlke, H., Horwath, W., 2020. Agricultural managed aquifer recharge — water quality factors to consider. *Calif. Agric.* 74(3), 144–154. <https://doi.org/10.3733/ca.2020a0020>.

Waterhouse, H., Arora, B., Spycher, N.F., Nico, P.S., Ulrich, C., Dahlke, H.E., Horwath, W.R., 2021. Influence of agricultural managed aquifer recharge (AgMAR) and stratigraphic heterogeneities on nitrate reduction in the deep subsurface. *Water Resour. Res.* 57, e2020WR029148. <https://doi.org/10.1029/2020WR029148>

Yadav, S.N., 1997, Formulation and Estimation of Nitrate-Nitrogen Leaching from Corn Cultivation. *J. Environ. Qual.* 26: 808-814.
<https://doi.org/10.2134/jeq1997.00472425002600030031x>

L. Appendix

Table 1. 37-year climate summary for five California Irrigation Management System stations used to input climate data for Root Zone Water Quality Model (RZWQM2) simulations. See Methods Section 2.2 for quality-control check, error-correction, and gap-filling procedures. Water balance components were summarized by water year (Oct-Sept) with evaporation and transpiration summarized from RZWQM2 business-as-usual median runs among all soils by station.

Parameter	Units	Location of CIMIS station				
		Durham	Davis	Parlier	Five Points	Shafter
Wind	m s ⁻¹	1.85	2.55	1.65	2.51	1.64
Relative Humidity	%	65.6	62.6	64.9	56.8	61.8
Insolation	kWh m ⁻²	4.82	4.96	4.94	5.19	5.27
Jan. Temp	°C	8.0	8.0	8.4	8.3	8.4
March Temp	°C	12.4	12.6	13.5	13.4	13.7
Mean Annual Temp	°C	16.1	16.0	17.1	17.0	17.1
July Temp	°C	24.4	23.5	26.5	25.9	26.4
Dormant season E _d	mm yr ⁻¹	189	165	131	114	101
Growing season E _g	mm yr ⁻¹	242	260	255	289	244
Crop ET	mm yr ⁻¹	502	667	605	758	593
Precip. minimum	mm yr ⁻¹	298	221	127	76	60
Precip. Median	mm yr ⁻¹	540	404	279	193	143
MAP	mm yr ⁻¹	569	444	280	211	159
P10	mm yr ⁻¹	648	518	345	278	182
Pmax	mm yr ⁻¹	1099	781	557	400	411
Pdorm	%	82.5	87.7	84.0	83.9	84.7
K-G arid	%	1.77	1.39	0.82	0.62	0.46

wind=mean annual; RH=relative humidity, mean annual; Insol=insolation, mean annual; Jan=January, mean; Mar=March, mean; MAT=mean annual temperature; Jul=July, mean; E_d=evaporation, dormant

season (Oct-Mar), median; E_g =evaporation, growing season (Apr-Sep), median; T_c =crop transpiration, median; P_{min} =minimum precipitation; P_{med} =median precipitation; MAP=mean annual precipitation; P_{10} =10th wettest water year precipitation ($\approx Q3$); P_{max} =maximum precipitation; P_{dorm} =dormant season (Oct-Mar) precipitation as annual percentage; $K-G_{arid}$ =Köppen-Geiger climate definition of aridity for climates where $>70\%$ of MAP falls during the winter months (Oct-Mar for northern hemisphere): $MAP / (MAT * 20)$, where MAP is in mm and MAT is in °C. If resulting aridity metric is <0.5 , the climate is arid desert; 0.5-1 is arid steppe—sometimes referred to as semi-arid.




Figure 5. Typical soil profile NO₃- (top row) and total NO₃- leached (bottom row) for a **coarse** (taxonomic family particle-size class) soil, comparing business-as-usual (no Flood-MAR) versus managed groundwater recharge (Flood-MAR(21d) = 60 cm yr⁻¹ across four applications on 21-day intervals), practiced during the 10 wettest winters from Davis (left column), the 2nd wettest climate modeled, thru Shafter (right column), the driest climate modeled. Note the difference in Y-axis scale in wet vs drier sites.




Figure 6. Typical soil profile NO_3^- (top row) and total NO_3^- leached (bottom row) for a **loamy** (taxonomic family particle-size class) soil, comparing business-as-usual (no Flood-MAR) versus managed groundwater recharge (Flood-MAR(21d) = 60 cm yr⁻¹ across four applications on 21-day intervals), practiced during the 10 wettest winters from Davis (left column), the 2nd wettest climate modeled, thru Shafter (right column), the driest climate modeled. Note the difference in Y-axis scale in wet vs drier sites.




Figure 7. Typical soil profile NO_3^- (top row) and total NO_3^- leached (bottom row) for a **fine** (taxonomic family particle-size class) soil, comparing business-as-usual (no Flood-MAR) versus managed groundwater recharge ((Flood-MAR(21d) = 60 cm yr⁻¹ across four applications on 21-day intervals), practiced during the 10 wettest winters in Davis (left column), the 2nd wettest climate modeled, thru Shafter (right column), the driest climate modeled. Note the difference in Y-axis scale in wet vs drier sites.

M. Factsheet/Database Template

1. **Project Title:** Techniques to minimize nitrate loss from the root zone during managed aquifer recharge
2. **Grant Agreement Number:** 20-1019
3. **Project Leaders:** Anthony Toby O'Geen and Helen Dahlke
4. **Start Year/End Year:** Start Date: 01/01/2021 End Date: 12/31/2025
5. **Location:** Statewide cropland
6. **County:** statewide cropland
7. **Highlights**

- Modeling suggests NO₃⁻ leaching risk during Managed Aquifer Recharge (Flood-MAR) varies by texture and climate
- NO₃⁻ leaching risk from Flood-MAR was low in locations with annual precipitation > 400 mm yr⁻¹
- NO₃⁻ leaching risk from Flood-MAR was elevated in locations with annual precipitation < 400 mm yr⁻¹
- In more arid locations NO₃⁻ leaching risk from Flood-MAR was highest for loamy soils
- Flood-MAR should be practiced with care in arid climates and especially after prolonged droughts
- Flood-MAR timing strategies (early vs late season), and water application frequency (3 vs. 7 vs. 21-day intervals) were negligible

8. Introduction

Agricultural management of floodwaters (Flood-MAR) is of broad interest in California as a tool to recharge aquifers. There are concerns to be analyzed before this practice can be safely implemented, such as contamination of groundwater by leaching soil nitrate (NO₃). This modeling exercise hypothesized that Flood-MAR will enhance NO₃ leaching vs. no Flood-MAR (business-as-usual). We hypothesized early Flood-MAR timing will leach less NO₃ than late Flood-MAR timing, due to lower rates of mineralization when soils are cooler. Mineralization generates NO₃ from soil organic matter decay. Additionally, frequency of Flood-MAR pulses (shorter interval between water applications) may leach less NO₃, due to less time for mineralization between Flood-MAR applications. Finally, NO₃ leaching risk is offset partially by denitrification in finer textured soils with longer periods of saturation and anaerobic conditions.


9. Methods/Management

This research evaluated contrasting seasonal timing and frequency of Flood-MAR as strategies to minimize NO₃ leaching by leveraging the Root Zone Water Quality Model (RZWQM), a widely validated tool developed and maintained by a team at USDA-ARS, to evaluate interaction of Flood-MAR with the N-cycle and inherent soil properties. Findings were tested in the field on semi-arid, coarse textured soils through an evaluation of the effect of cover crops on NO₃ leaching during Flood-MAR.

RZWQM modeling included 33 Central Valley soils representing distinct particle-size classes (termed here Coarse, Loamy and Fine) and five different 37-year climate records obtained from CIMIS, spanning a precipitation gradient from relatively wet-to-dry in space and time (n=990 unique scenarios). The climatic gradient summarized by town from wettest to driest is as follows: Durham, Davis, Parlier, Five Points, and Shafter. Simulations extended to a depth of 150 cm. Leaching values represent the NO₃

flux at the bottom boundary of this simulation domain. Biogeochemical and physical parameters were established using end-of-run values from a preliminary 37-year business-as-usual run of each unique soil x climate modeling combination (n=165). This produced unique initial biogeochemical conditions for each of the soil x climate combinations to test again under another 37-year business-as-usual run and contrasting Flood-MAR strategies. In simulations, Flood-MAR was practiced during the 10-wettest water years of each 37-year climate record, applying 600-cm additional water via Flood-MAR. In a Flood-MAR year, four 15-cm water applications were made in either January or March, using a frequency of either 3- or 7-day intervals. A fifth scenario tested a 21-day Flood-MAR interval January-March.

10. Findings



Multi-decadal RZWQM simulations suggest Flood-MAR can be used with near negligible risk of additional NO_3 leaching in relatively wet Central Valley locations (Durham and Davis, median annual precipitation $> 400 \text{ mm yr}^{-1}$) across a range of soil textures. Steady-state residual NO_3 in the wetter climates (Durham and Davis) were typically $60\text{-}100 \text{ kg N ha}^{-1}$ after 37-years of the business-as-usual scenario (Fig. 1).

This is because in-situ precipitation during the wet years, when Flood-




Figure 2. Effect on 37-year, cumulative nitrate leached of different Flood-MAR strategies, including 3-day (d) and 7-d frequency events practiced either in January (Jan) or March (Mar) or 21-d frequency events spanning early Jan thru late Mar. Additional nitrate leached is relative to each soil's business-as-usual run (no Flood-MAR). Each plot is grouped by three generalized taxonomic family particle-size classes: coarse; loamy; and fine (n=11 soils per class); and shows the different climates modeled from wettest (Durham) to driest (Shafter). See Table 1 for descriptions of climates

MAR is expected to be practiced, removed most residual NO_3 through deep percolation. This is true even in the finest textured soils, which are most difficult to leach due to high microporosity. As precipitation declines, the Flood-MAR NO_3 leaching risk increased most clearly in loamy soils, even though the central tendency did not differ substantially across textural groups (Fig. 1 & 2). Additional nitrate leaching risk increased in dry climates, because lack of precipitation allowed for residual NO_3 accumulation across growing seasons.

Loamy soils tended to present the greatest possibility of risk of additional NO_3 leaching with Flood-MAR in drier climates (Figure 2). In the driest climate (Shafter), 4 of 11 loamy soils leached $>3000 \text{ kg additional NO}_3\text{-N ha}^{-1}$ using 21-day frequency Flood-MAR with median fluxes of $1,270 \text{ kg additional NO}_3\text{-N ha}^{-1}$. In fine soils, NO_3 leaching risk was mitigated by denitrification, preventing build-up of residual NO_3 . Flood-MAR timing strategies (January Flood-MAR vs. March Flood-MAR, combined with variable pauses among applications (3 vs. 7 vs. 21-day intervals, the latter January-March Flood-MAR) had only a negligible effect on NO_3 leaching risk. In fact, the effect of Flood-MAR timing strategies was only noticeable in wet climates where additional NO_3 leaching risk was comparably very low. While results demonstrated that Flood-MAR practices would be expected to increase net NO_3 flux to groundwater across all climates and soils, consistent Flood-MAR practices would also be expected to improve groundwater quality compared to business-as-usual irrigated agriculture. This is due to sustained provision of higher quality deep percolation water, which is especially limited in dry climates. Thus, climates with median precipitation $> 400 \text{ mm yr}^{-1}$ were sufficient to leach rootzones in this simulated, fertilized agroecosystem, suggesting that Flood-MAR practiced in wetter climates is of low additional NO_3 leaching risk (Fig. 2).




Figure 3. Mean nitrate concentration before and after Flood-MAR in soils near Modesto and Shafter.

support modeled outcomes. For example, both field trials showed reductions in profile NO₃ concentration after Flood-MAR (Fig. 3). Loamy soils in dry (Modesto) and relatively moist settings (Durham) had high residual nitrate near the soil surface, but values were identical after Flood-MAR possibly due to mineralization of organic matter.

In summary, we found that residual soil nitrate was highest in the south where winter precipitation was too low to completely flush soil profiles. We also found that loamy soils had higher residual soil nitrate because these soils have high water retention, thus had higher nitrate leaching risk. Fine textured soils also have high water retention, but the water filled pore space in these soils remained high enough to promote denitrification resulting in a gaseous loss of nitrate, lowering leaching risks. Modeling different timings and durations of flood-MAR showed no meaningful effect on additional nitrate leaching.

The most direct mechanistic explanation for additional nitrate leaching risk in loamy soils from drier climates is due to their high capacity to retain water and NO₃. Loamy soils require more percolating water to leach effectively compared to coarse soils, explaining their conduciveness to residual NO₃ accumulation. Although coarse soils typically present the greatest risk to NO₃ leaching in agriculture, this truism did not hold up to evaluations of the effect of Flood-MAR on additional NO₃ leaching risk. Except in the driest climates, precipitation is sufficient to leach residual NO₃, such that the additional NO₃ leaching risk from Flood-MAR is typically lower in coarse soils compared to loamy soils.

Field Flood-MAR trials from almond fields sampled at dry (Modesto) and moist (Durham) sites generally

N. Copy of product/result

We have two products to share. A peer reviewed scientific journal article, and an extension bulletin. The draft of the extension bulletin can be found below. The publication is also included at the end of the report.

Understanding nitrate leaching risk for managed aquifer recharge in agriculture

A.T. O'Geen, C. Kisliuk, and H. Dahlke

Introduction

As climate change and chronic groundwater overdraft intensify water scarcity in California, managed aquifer recharge in agriculture (Ag-MAR) is gaining popularity as a strategy to replenish depleted groundwater supplies (O'Geen et al., 2015; Dahlke et al., 2018; Bachand et al., 2022; Levintal, et al., 2023). Ag-MAR involves the intentional application of surface water on agricultural lands to recharge aquifers, providing temporary storage for excess floodwater during wet periods and augmenting groundwater availability during drought. In addition to groundwater replenishment, Ag-MAR offers a range of co-benefits, including flood risk mitigation, land subsidence abatement, wetland restoration, and enhanced ecosystem functions such as habitat support for migratory birds and salmonids (Damigos et al., 2017).

Ag-MAR is a form of flood-managed aquifer recharge (Flood-MAR) in which surface water is intentionally applied to croplands, low-lying and nearly level rangelands, or fallow farmland. Ag-MAR is distinct from other Flood-MAR approaches because it involves active agricultural production systems, where recharge can pose risks of crop damage and potential groundwater contamination from fertilizers and pesticides used in agricultural production. Ag-MAR typically has shorter water application durations, and it is not always applied to soils with high-infiltration compared to other types of groundwater banking practices.

California cropland is ideal for ground water banking due to its vast extent. Of the 17.5 million acres of irrigated and non-irrigated farmland, according to the Soil Agricultural Groundwater Banking Index (SAGBI), over 5 million acres were ranked excellent, good, or moderately good for this practice (O'Geen et al., 2015). An additional 12.6 million acres of less suitable land was identified by SAGBI. Much of this land is considered poorly suited to Ag-MAR because soils are slowly permeable to water. Prolonged soil saturation can harm certain crops, especially perennial crops. Principal among the risks of widespread Ag-MAR adoption is nitrate leaching.

Nitrate (NO_3^-) leaching to groundwater is a major public-health concern in agricultural regions across California. Nitrate enters groundwater primarily through downward leaching of excess nitrogen that accumulates in soils and the vadose zone as a result of

inefficient fertilizer management. California's cropland is responsible for 96% of human-generated NO_3^- pollution to groundwater (Harter et al., 2012). Consumption of water with more than 10 ppm NO_3^- -N, the maximum contaminant level (MCL) in California, is believed to cause blue baby syndrome, cancer, and birth defects. Ransom & Harter (2017) found that a small number of private wells (usually shallow) will surpass the maximum contaminant level (MCL = 10 ppm nitrate-nitrogen), and that a large part of the Central Valley will have elevated NO_3^- (above 5 ppm). Deeper public wells are less threatened, but many of them are predicted to have elevated nitrate levels, too. When considering adoption of flooding farmland for groundwater recharge, stakeholders should evaluate the NO_3^- leaching risk—caused by legacy NO_3^- . This risk is mediated by the nitrogen cycle (i.e. nitrogen transformation processes including mineralization, nitrification and denitrification) and the water balance of an Ag-MAR activity.

The Nitrogen Cycle during Ag-MAR

The fate of N in soil is controlled by the N-cycle which encompasses additions, losses and transformations of N within soil (Fig. 1). For the most part, the N-cycle is mediated by microbes, but environmental conditions such as climate, soil type and management play a role in the fate of N, especially in the context of Ag-MAR timing, duration and amount of water applied (Levintal et al., 2023; Cui et al., 2025).

Additions and losses of N go hand-in-hand when considering NO_3^- contamination risk during Ag-MAR. Additions of N arise mainly from fertilizer application and to a lesser extent N fixation by plants and microbes. Losses of soil N are mainly a result of removal via harvest and leaching but also include gaseous losses of volatile N forms as ammonia volatilization and denitrification. High residual NO_3^- concentration in soil after harvest is the main contributor to NO_3^- leaching risk during Ag-MAR. Modeling suggests the contribution of residual NO_3^- to deep percolation during Ag-MAR is highest in more arid portions of the State such as the southern San Joaquin Valley where precipitation is low and residual NO_3^- build-up is greater. These conditions allow NO_3^- to persist in soil over fall and winter months (Devine et al., 2025). The best strategy to reduce NO_3^- leaching risk during Ag-MAR is to achieve low residual NO_3^- prior to flooding.

Transformations in the N-cycle affect the form of N in soil, which in turn influence N leaching risk. For example, plant uptake converts inorganic N (NH_4^+ and NO_3^-) into organic N that is incorporated into plant biomass and residues. As a result, cover crops can be an effective strategy for scavenging residual soil N and reducing leaching risk during Ag-MAR, provided that crops can be established early enough in the season. Organic N contained in crop residues and soil organic matter can be converted back into inorganic ammonium (NH_4^+) through microbial decomposition, a process known as mineralization (Fig 1). Ammonium poses a low risk of leaching because it is positively charged and attracted to negatively charged clay minerals. However, ammonium is rapidly transformed to NO_3^- in well drained soils through the microbial process of nitrification. Because nitrate is negatively charged, it is highly mobile in soil solution and therefore susceptible to leaching especially during periods of excess water application

such as Ag-MAR. In cool or dry climates mineralization rates are strongly reduced during much of the off-season due to low soil temperature, low soil moisture, or both (Geisseler et al., 2019) indicating a relatively low contribution to nitrate leaching risk during Ag-MAR. Ag-MAR events may stimulate mineralization where microbial activity is limited by soil moisture, potentially increasing nitrate availability during or following recharge events (Murphy et al., 2021).

Like with plant uptake, both ammonium and NO_3^- can be temporarily immobilized by microbial uptake. Immobilization occurs when large additions of carbon-rich materials are added to soil which stimulates microbial activity. When immobilized, N is not available for plant uptake or at risk of leaching losses. This process would be difficult to manage/optimize for Ag-MAR, given that immobilization is a temporary process governed by microbial activity.




Figure 1. The fate of N in the nitrogen cycle.

When soil become saturated and oxygen availability is low, NO_3^- can be reduced by denitrifying microbes to dinitrogen (N_2) gas, resulting in a loss of nitrogen to the atmosphere. Modeling studies have shown that denitrification accounts for a small but meaningful fraction of soil nitrate removal during Ag-MAR, generally contributing less than 25% of total NO_3^- losses, depending on soil type and environmental conditions (Cui et al., 2025). Other studies have shown substantially higher denitrification rates in fine-

textured soils under simulated Ag-MAR conditions. This enhanced denitrification is believed to be large enough to reduce nitrate leaching risk in fine-textured soils compared to loamy and coarse-textured soils, particularly where residual nitrate levels are high (Devine et al., 2025). The extent of denitrification during Ag-MAR largely depends on the duration of saturated conditions, which in turn depends on both water application duration and soil texture. Secondary controls include soil temperature and the availability of organic carbon as electron donor for denitrifying microbes (Braker et al., 2010).

Managing Soil Nitrate Concentration in Deep Percolation from Ag-MAR

The greatest risk of N leaching occurs when soil NO_3^- levels are high. The most important step in reducing groundwater contamination during Ag-MAR is to reduce the concentration of NO_3^- when leaching occurs. The first step to minimize NO_3^- leaching is to know the residual NO_3^- concentrations in soil prior to Ag-MAR. A soil test in fall or early winter should be considered. Residual soil NO_3^- tests are also a valuable post-harvest evaluation to inform N management practices. The soil test should be representative of the entire soil profile (upper 5 feet). Soil testing at times closer to an Ag-MAR event is better to limit the potential changes associated with N cycling discussed above. No threshold level of residual soil NO_3^- exists because concentrations are dynamic, changing with water content and due to N-cycling processes.

It is critical that sufficient water is applied during Ag-MAR to ensure that NO_3^- concentrations in deep percolation remain below the drinking water MCL. Table 1 provides mass balance estimates of the Ag-MAR water volumes needed to dilute NO_3^- concentration in deep percolation to below the drinking water standard (≤ 10 ppm), accounting for a range of residual soil NO_3^- concentration and soil textures. The results show that substantially less applied water is needed to meet the MCL when residual soil NO_3^- concentrations are low (< 20 ppm). For example, at a residual soil NO_3^- concentration of 20 ppm, approximately 2.7 feet of applied water is required to dilute pore water nitrate concentrations below the MCL in fine textured soils, compared to only 0.7 ft in coarse-textured soil. Fine-textured soils require more water compared to coarse textured soils because they retain more NO_3^- per unit depth (Levintal, et al., 2023). At higher residual nitrate levels, required recharge volumes increase substantially. For coarse-textured soils with residual NO_3^- concentrations of 80 ppm, at least 3.6 ft of applied water is needed to reach the MCL in deep percolation during Ag-MAR, whereas fine-textured soils with the same residual NO_3^- concentration require approximately 10.7 feet of applied water (Table 1). These estimates assume that NO_3^- concentrations in the source water are negligible.

Table 1. Estimates of the amount of Ag-MAR water needed to reduce nitrate concentration in deep percolation to levels below the maximum contaminant level (MCL = 10 ppm) considering different residual soil nitrate levels and soil textures.

Residual Soil Nitrate (ppm)	Coarse texture	Fine texture
	Amount of water needed ----- (ft) -----	
15	0.7	2.0
20	0.9	2.7
30	1.3	4.0
40	1.8	5.3
50	2.2	6.7
60	2.7	8.0
80	3.6	10.7
100	4.4	13.3

Estimates are calculated for a total soil depth of 4 feet.
Assumes nitrate concentration in Ag-MAR water is below MCL.

Field monitoring before and after Ag-MAR demonstrates the dilution effect on soil NO_3^- concentration. In uniformly sandy soil, variable NO_3^- concentrations with depth were reduced to near zero after Ag-MAR (Fig. 2). In contrast, soils with clay-rich layers appear to inhibit complete flushing of NO_3^- after Ag-MAR (Fig. 3) and NO_3^- transport overall is slower. The peak in NO_3^- concentration at depth after Ag-MAR in Figure 3 suggests that not enough water was applied to fully leach the deep profile, or a water restrictive layer exists (note the large clay increase at 1000 cm depth), which may prevent deep percolation beyond that point.




Figure 2. Soil nitrate concentration before and after deep percolation (a) in a uniformly coarse textured soil (b).

Modeling Predictions

Ag-MAR has the potential to modify nitrogen cycling in ways that may benefit groundwater quality. Modeling studies have evaluated how adjusting the applied water amount, timing, duration, and frequency of recharge events can optimize key nitrogen transformation processes. For example, one study tested four 15-cm Ag-MAR water applications in either January or March, with application intervals of 3 or 7 days. A fifth scenario applied recharge from January through March at 21-day intervals. These simulations showed no significant difference in nitrate leaching as a function of timing or water application frequency (Devine et al., 2025). A separate modeling study found that intermittent Ag-MAR events allowing drainage between applications stimulated soil nitrogen mineralization, resulting in greater NO_3^- leaching during subsequent recharge events (Cui et al., 2025). Continuous flooding, with recharge durations ranging from 8 to 20 days, produced the lowest NO_3^- leaching losses. Finally, implementing Ag-MAR closer to the growing season increased the availability of both water and nitrogen for crop uptake.




Figure 3. Soil nitrate concentration before and after deep percolation (a) in a soil with contrasting texture in the deeper profile (b).

Implications

If Ag-MAR becomes a widespread and sustained practice, it has the potential to progressively dilute NO_3^- concentrations in soils and sediments, ultimately leading to improved groundwater quality over time. However, these benefits are not automatic and can only be realized if nitrate dynamics are explicitly considered during Ag-MAR planning and implementation. Achieving long-term groundwater quality improvements will also depend on continued adoption of careful nutrient management during the growing season to minimize residual NO_3^- following harvest.

Online Tools

UC Davis's California Soil Resources Laboratory developed the [Soil Agricultural Groundwater Banking Index](https://soilmap2-1.lawr.ucdavis.edu/root-zone/), a map that identifies groundwater recharge suitability geographically based on factors including deep percolation, root zone residence time <https://soilmap2-1.lawr.ucdavis.edu/root-zone/>, topography, chemical limitations, and soil surface condition. It is available at <casoilresource.lawr.ucdavis.edu/sagbi/>.

The Nitrogen Leaching Hazard Index was developed to predict nitrate leaching risk based on customizable inputs including crop type, soil type, and irrigation method. It is available at <https://soilmap4-1.lawr.ucdavis.edu/nitrogen-hazard-index/>

UC Davis and the CDFA collaborated on the [California Fertilization Guidelines](https://geisseler.ucdavis.edu/Guidelines/Home.html), that help predict crop fertilization requirements for the purposes of avoiding overfertilization and nitrate pollution. It is available at geisseler.ucdavis.edu/Guidelines/Home.html.

The *time-to-trafficability* tool (<https://soilmap4-1.lawr.ucdavis.edu/soil-trafficability/>) is intended to help California growers identify when fields are generally *trafficable* after *deep soil wetting* during crop dormancy or winter fallow periods. The tool applies to wetting situations such as managed aquifer recharge projects and large rain or flood events. The objective of the tool is to help growers avoid physical soil damage by agricultural vehicles, so estimates are relatively conservative.

References

- Bachand, P., S.Roy, D. Stern, J. Choperena, D. Cameron, and W. Horwath. 2016. On-farm flood capture could reduce groundwater overdraft in Kings River Basin. *California Agriculture*, Volume 70: 200-207. <https://doi.org/10.3733/ca.2016a0018>
- Braker, G., J. Schwarz, and R. Conrad. 2010. Influence of temperature on the composition and activity of denitrifying soil communities. *FEMS Microbiology Ecology*.

Volume 73, Issue 1, July 2010, Pages 134–148. <https://doi.org/10.1111/j.1574-6941.2010.00884.x>

Cui, W. T. Zhou, E. Levintal, C. Prieto Garcia, I. Kisekka, H.E. Dahlke. 2025 Modeling the impact of agricultural managed aquifer recharge (Ag-MAR) on soil water and nitrogen dynamics of the growing season. *Agricultural Water Management* 317 <https://doi.org/10.1016/j.agwat.2025.109623>

Dahlke, H.E., A.G. Brown, S. Orloff, D. Putnam and T. O'Geen, 2018. Managed winter flooding of alfalfa recharges groundwater with minimal crop damage. *California Agriculture Journal*. <https://doi.org/10.3733/ca.2018a0001>

Damigos, D., Tentes, G., Balzarini, M., Furlanis, F., & Vianello, A. (2017). Revealing the economic value of managed aquifer recharge: Evidence from a contingent valuation study in Italy. *Water Resources Research*, 53(8), 6597–6611. <https://doi.org/10.1002/2016WR020281>

Devine, S., Dahlke, H., Harter, T., Kisekka, I., Najm, M. A., & O'Geen, A. T. (2025). Modeling nitrate leaching risk from flood managed aquifer recharge in California's agricultural lands. *Vadose Zone Journal*, 24(6), e70058.

Harter, T., J. R. Lund, J. Darby, G. E. Fogg, R. Howitt, K. K. Jessoe, G. S. Pettygrove, J. F. Quinn, J. H. Viers, D. B. Boyle, H. E. Canada, N. DeLaMora, K. N. Dzurella, A. Fryjoff-Hung, A. D. Hollander, K. L. Honeycutt, M. W. Jenkins, V. B. Jensen, A. M. King, G. Kourakos, D. Liptzin, E. M. Lopez, M. M. Mayzelle, A. McNally, J. Medellin Azuara, and T. S. Rosenstock. 2012. Addressing Nitrate in California's Drinking Water with a Focus on Tulare Lake Basin and Salinas Valley Groundwater. Report for the State Water Resources Control Board Report to the Legislature. Center for Watershed Sciences, University of California, Davis. <https://ucanr.edu/sites/groundwaternitrate/files/138956.pdf>

Levintal, E., L. Huang, C.P. Garcia, A. Coyotl, M.W. Fidelibus, W.R. Horwath, J.L.M. Rodrigues, and H. Dahlke. 2022. Nitrogen fate during agricultural managed aquifer recharge: Linking plant response, hydrologic, and geochemical processes. *Science of the Total Environment*, 864 (2023). <https://doi.org/10.1016/j.scitotenv.2022.161206>

O'Geen, A. , M. Saal, H. Dahlke, D. Doll, R. Elkins, A. Fulton, G. Fogg, T. Harter, J. Hopmans, C. Ingels, F. Niederholzer, S. Sandoval, P. Verdegaal, and M. Walkinshaw. 2015. Soil suitability index identifies potential areas for groundwater banking on agricultural lands. *California Agriculture* 69(2):75-84. <https://doi.org/10.3733/ca.v069n02p75>

Ransom, K. and T. Harter. 2017. Groundwater nitrate sources and contamination in the Central Valley. Center for Watershed Sciences, University of California, Davis. <https://californiawaterblog.com/2017/09/17/groundwater-nitrate-sources-and-contamination-in-the-central-valley>

ORIGINAL ARTICLE

Modeling nitrate leaching risk from flood managed aquifer recharge in California's agricultural lands

Scott Devine  | Helen Dahlke  | Thomas Harter | Isaya Kisekka  |
Majdi Abou Najm  | Anthony T. O'Geen 

Department of Land, Air, and Water Resources, University of California Davis, Davis, California, USA

Correspondence

Anthony T. O'Geen, Department of Land, Air, and Water Resources, University of California Davis, Davis, CA 95616-8627, USA.

Email: atogeen@ucdavis.edu

Assigned to Associate Editor Shengmin Luo.

Funding information

California Department of Food and Agriculture—Fertilizer Research and Education Program

Abstract

Flood managed aquifer recharge (Flood-MAR) is an emerging practice to enhance groundwater sustainability through recharge of aquifers. However, in agricultural systems Flood-MAR may harm groundwater quality, given residual soil nitrate (NO_3^-). This research evaluated Flood-MAR NO_3^- leaching risk across a precipitation and soil textural gradient in a globally important agricultural region, California's Central Valley, and whether Flood-MAR timing strategies could mitigate the risk. Using multi-decadal root zone water quality model simulations of irrigated and fertilized maize ($250 \text{ kg N ha}^{-1} \text{ year}^{-1}$) on well-drained soils as a representative case-study, results suggest Flood-MAR can be used with near-negligible additional NO_3^- leaching in locations with median annual precipitation $>400 \text{ mm year}^{-1}$. In those locations, wet-year precipitation leached most residual NO_3^- without practicing Flood-MAR. At drier locations, Flood-MAR NO_3^- leaching risk increased most clearly in loamy soils. Additional NO_3^- leaching risk increased in drier climates because minimal precipitation-driven deep percolation maintained residual NO_3^- accumulation across growing seasons. In fine-texture soils, NO_3^- leaching risk was mitigated by denitrification, preventing residual NO_3^- accumulation. Flood-MAR practices diminished denitrification, leaching NO_3^- rapidly when soils were colder and biogeochemically inactive, and thereby decreased growing season denitrification when soils were warmer and denitrification rates higher. Effects of Flood-MAR timing strategies (January Flood-MAR vs. March Flood-MAR), combined with variable pauses among applications (3- vs. 7- vs. 21-day intervals) were negligible. Infrequent Flood-MAR should be practiced with care in arid climates and especially after prolonged droughts coinciding with more limited irrigation water supplies that constrain salt-leaching practices all favoring residual NO_3^- accumulation.

Abbreviations: BAU, business-as-usual; CERES-Maize, crop environment resource synthesis model; CIMIS, California Irrigation Management Information System; Flood-MAR, Flood Managed Aquifer Recharge; MCL, maximum contaminant level; PRISM, parameter-elevation regressions on independent slopes model; RZWQM2, root zone water quality model; SIC, simulated initial conditions; SOM, soil organic matter; SSURGO, Soil Survey Geographic database; UAN, urea ammonium nitrate.

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2025 The Author(s). *Vadose Zone Journal* published by Wiley Periodicals LLC on behalf of Soil Science Society of America.

Plain Language Summary

Agricultural management of floodwater (Flood-MAR) is a new practice where floodwaters are applied to agricultural fields to recharge groundwater. Flood-MAR may harm groundwater quality by leaching soil nitrate (NO_3^-) into groundwater. This modeling study evaluated the Flood-MAR NO_3^- leaching risk in different climates and soil textures in California. It evaluated whether Flood-MAR timing strategies (early- vs. late-season irrigation application strategies) influenced risk. Flood-MAR had near-negligible NO_3^- leaching risk in locations with rainfall >400 mm year $^{-1}$. At drier locations, Flood-MAR NO_3^- leaching risk was highest (especially in loamy soils) because low rainfall allowed NO_3^- to build up in soils from year to year. Different Flood-MAR timing strategies (early season vs. late season), combined with variable pauses in water applications (3- vs. 7- vs. 21-day intervals), showed no difference in NO_3^- leaching. Flood-MAR should be practiced with care in arid climates.

1 | INTRODUCTION

Globally, irrigated landscapes have more than doubled in extent since the 1960s to 342 million ha in 2019, helping boost agricultural production at a rate faster than the total area under cultivation (FAO, 2021). However, when practiced in semiarid and arid climates with a natural climatic water deficit, irrigated agriculture requires enormous freshwater withdrawals to meet irrigated crop evapotranspiration demands; for example, in California, 81% of groundwater use has been attributed to irrigated agriculture (Nelson, 2012). This intense reliance on groundwater has led to numerous occurrences where groundwater extraction rates greatly exceed recharge rates underlying irrigated regions (Herbert & Döll, 2019; Siebert et al., 2010). In California, where globally important agriculture has revenues now exceeding \$60 billion annually, multiyear droughts in recent decades have highlighted groundwater depletion as an existential threat to the long-term viability of irrigation and even access to groundwater in overlying communities (Liu et al., 2022).

The agricultural management of floodwaters (Flood-MAR) is of broad interest as a tool to recharge aquifers, offset groundwater pumping, and mitigate downstream flood risk (CDWR, 2018). This practice involves flooding agricultural fields during the winter and spring, and in some cases early summer, when rivers can reach high magnitude flows during wet years as a result of atmospheric river events and snowmelt (Kocis & Dahlke, 2017). Novel approaches are necessary to sustain irrigated agriculture in the face of multiple challenges. These include new public policy constraints on groundwater use, such as the Sustainable Groundwater Management Act requiring groundwater sustainability plans in overdrafted basins (CDWR, 2022), and additional challenges imposed by climate change, such as more intense precipitation whiplash

(Swain et al., 2018) and decreased mountain snowpack, a natural reservoir for California's irrigated agriculture that will diminish with warming (Qin et al., 2020).

However, there are concerns to be analyzed before Flood-MAR can be safely implemented, such as contamination of groundwater by flushing out residual soil nitrate (NO_3^-) during crop dormancy or fallowing at times that coincide with high magnitude river flows. Our research assumed an overarching hypothesis that Flood-MAR would enhance NO_3^- leaching compared to no Flood-MAR (business-as-usual [BAU]), following the concern that Flood-MAR would mobilize residual NO_3^- in fertilized agroecosystems (Levinthal et al., 2022). For example, in relatively deep vadose zone samples under California citrus orchard fertilizer trials, Harter et al. (2005) reported residual nitrate levels of 218–477 kg NO_3^- -N ha $^{-1}$ in 16-m deep cores. Bachand et al. (2014) measured residual NO_3^- in western Fresno County fields (San Joaquin Valley, California) where Flood-MAR had been practiced for one season and observed a residual NO_3^- quantity less than 15%, on average, of the soil profile NO_3^- (0–3 m) in wine grape control fields. In a deep soil coring study (0–30 m) across 36 study sites (12 fields) in the same region, Waterhouse et al. (2020) found relatively high residual NO_3^- across tomatoes and almonds (737–966 and 420–1240 kg NO_3^- -N ha $^{-1}$, respectively) with a lower range for wine grapes where some growers tend to apply less fertilizer (75–1371 kg NO_3^- -N ha $^{-1}$), but overall still demonstrating the risk for mobilizing residual NO_3^- with Flood-MAR.

We evaluated scenarios of contrasting seasonal timing and frequency of Flood-MAR as strategies to possibly minimize NO_3^- leaching risk by leveraging the root zone water quality model (RZWQM2), a widely validated tool developed and maintained by a team of USDA Agricultural Research Service scientists. We explored the primary controls (climate and soil

properties) influencing the interaction of Flood-MAR with the N-cycle. Laboratory-based soil column leaching experiments demonstrated that the frequency of wetting in the field can potentially influence NO_3^- leaching. For example, when 15-cm water pulses were applied every 1–2 weeks, total NO_3^- leached exceeded initial residual NO_3^- by 30%–50% because N was mineralized from the organic pool between pulses. When 15-cm water pulses were applied every 3 days, total NO_3^- leached was less than the initial residual NO_3^- (Murphy et al., 2021). Following this finding, we hypothesized that early-winter Flood-MAR timing would leach less NO_3^- than late-winter or early-spring Flood-MAR timing over the long term, due to lower rates of N mineralization when soils are cooler. Additionally, we hypothesized that higher frequency Flood-MAR pulses (shorter interval between water applications) would leach less NO_3^- , since there would be less time for soils to generate NO_3^- from mineralization of soil organic matter (SOM) between Flood-MAR applications. As a counteracting process to the effect of leaching, Waterhouse et al. (2021) demonstrated that prolonged Flood-MAR may enhance some NO_3^- removal via denitrification. Following this, we also hypothesized that Flood-MAR NO_3^- leaching risk would be partially offset by denitrification in finer textured soils, which would require more extended periods of soil saturation to achieve similar recharge depths compared to coarser soils, leading to anaerobic conditions. We expected this mitigation to be amplified when Flood-MAR was repeatedly practiced in the latter part of the wet season when soils are warmer and temperature-driven denitrification rates are potentially higher. RZWQM2 was used to simulate a multi-decadal BAU scenario and five different Flood-MAR approaches in the context of a fertilized ($250 \text{ kg N ha}^{-1} \text{ year}^{-1}$) and irrigated maize agroecosystem with winter fallows across different representative soils and climates, as described below.

2 | MATERIALS AND METHODS

2.1 | Overview of RZWQM2

RZWQM2 is a comprehensive process model that simulates fluxes of soil water and heat, organic matter cycling, and biogeochemical transport of N, linked to crop growth and nutrient and water uptake models, all paired to an agriculture management module (Ahuja et al., 2000). A primary use of RZWQM2 has been to calibrate and then predict the hydrologic response of crop–soil–climate combinations to various agricultural management systems, including effects on groundwater or surface water quality through runoff, leaching, or tile drainage (Kisekka et al., 2017; Ma et al., 2006). In California, for example, RZWQM2 has been used to model the application of dairy manure to cropland and its effects

Core Ideas

- Modeling suggests NO_3^- leaching risk during flood managed aquifer recharge (Flood-MAR) varies by texture and climate.
- NO_3^- leaching risk diminished from Flood-MAR in locations with annual precipitation $>400 \text{ mm year}^{-1}$.
- Flood-MAR should be practiced with care in arid climates and especially after prolonged droughts.
- Flood-MAR timing strategies (early vs. late season) and durations (3- vs. 7- vs. 21-day intervals) were negligible.

on various greenhouse gas emission components (Geisseler et al., 2012) and NO_3^- leaching to groundwater from an intensively monitored almond orchard site (Domagalski et al., 2008).

The RZWQM2 model embedded with DSSAT-CSM v4.0 (decision support systems for agrotechnology transfer cropping system model) CERES-Maize (crop environment resource synthesis model) crop growth module was used in this study (Ma et al., 2011). Infiltration and runoff are simulated by RZWQM2 using a modified form of the Green–Ampt equation (Green & Ampt, 1911). Unsaturated soil water flow and soil water redistribution are modeled using a one-dimensional (1D) Richards equation (Richards, 1931). Macropore flow is governed by the Poiseuille’s law. When rainfall or irrigation exceeds the soil’s infiltration capacity, RZWQM2 generates runoff. The potential evaporation and transpiration demand of the atmosphere were computed using the Shuttleworth–Wallace ET model (Ahuja et al., 2000). The Shuttleworth–Wallace ET model in RZWQM2 is an extension of the Penman–Monteith ET model, but the former considers incomplete canopy cover and plant height in potential evaporation and transpiration estimations (Kisekka et al., 2017). Actual root water uptake described by the sink term in Richards’s equation was calculated numerically using the Nimah and Hanks (1973) procedure. Actual root water uptake, potential evaporation, and potential transpiration were used in computing water stress factors used in CERES-Maize to modulate plant growth processes like leaf growth as soil water gets depleted. Carbon–nitrogen (N) cycles are driven by microbial biomass and environmental factors such as temperature and soil water content.

RZWQM2 simulations were performed on soil profiles (152- to 218-cm deep) with properties defined by horizon based on the Soil Survey Geographic database (SSURGO). RZWQM2 then automatically discretizes these user-defined soil horizons into multiple numerical nodes per horizon to simulate 1D soil water and heat transport (i.e., nodes placed

TABLE 2 Summary of conditions simulated using the root zone water quality model.

Scenario	Flood-MAR application frequency	Season
BAU (business-as-usual)	NA	NA
High frequency-late	15 cm H ₂ O applied every 3 days	March
High frequency-early	15 cm H ₂ O applied every 3 days	January
Low frequency-late	15 cm H ₂ O applied every 7 days	March
Low frequency-early	15 cm H ₂ O applied every 7 days	January
Extended frequency	15 cm H ₂ O applied every 21 days	January through March

Note: Each Flood-MAR scenario consisted of four applications of 15 cm of water performed during the 10 wettest years of the 37-year simulation period. BAU received no Flood-MAR treatment, thus information for application frequency and season was not applicable (NA).

This approach was used because soil survey data reflect soil properties of representative pedons from locations with natural conditions rather than irrigated fields, which have higher biomass inputs and a wetter moisture regime. Specifically, SOM pools were first parameterized by soil horizon using data from the SSURGO and initially assumed to be distributed across the SOM pools as follows: the slow-cycling pool was assumed to represent 80% of total SOM; the intermediate-cycling pool 15%; and the fast-cycling pool 5% across all horizons, following Ma et al. (1998) and Cameira et al. (2007), but then allowed to equilibrate during the 37-year SIC run, saving the end-of-run values as the SIC. This produced unique initial biogeochemical conditions for each of the soil × climate combinations to be tested again under another 37-year BAU run (without Flood-MAR treatments) and five 37-year model runs with contrasting Flood-MAR strategies, as described next.

In RZWQM2 simulations, Flood-MAR was practiced during the 10 wettest water years (October–September) of each specific 37-year climate record, applying 600-cm additional water (60 cm year⁻¹) through Flood-MAR. During a Flood-MAR year, four 15-cm water applications were made in either January or March, using an application frequency of either 3- or 7-day intervals (Table 2). A fifth scenario tested applications from January to March with 21-day intervals. For fine-textured soils with slower saturated hydraulic conductivities, Flood-MAR was practiced up to nine continuous days to achieve 15-cm applied water in each event, depending on the soil's depth-weighted hydraulic conductivity, using 3-, 7-, or 21-day periods between these continuous periods of Flood-MAR, depending on the scenario. RZWQM2 can mimic surface ponding by adjusting the maximum surface storage parameter. In this study, in order to realistically represent FloodMAR, time periods to achieve 15-cm infiltrated water had to be determined a priori as shown in Table 2.

2.1.2 | Simulated cropping system details

The DSSAT-CSM v4.0 CERES-Maize model was used to represent crop growth and development in the RZWQM2

model (Ma et al., 2006) as continuous maize with winter fallow. The CERES-Maize model is a radiation-based mechanistic crop model that predicts maize growth and development based on weather (precipitation, solar radiation, maximum and minimum temperature, and to a lesser extent photoperiod), management practices (plant population, row spacing, planting date, and irrigation management), and six cultivar coefficients (P1, P2, P5, G2, G3, and PHINT) as described in Hoogenboom et al. (2019). More details about the RZWQM2 can be found in Ahuja et al. (2000).

The objective for the project here was to have a stable crop growth model with reliable, consistent plant growth and N accumulation each year across all soil × climate scenarios, accepting that some minor differences would be expected. Ideally, similar yields and thus N removal through harvest would result in similar potentials for residual nitrate buildup due to less removal than applied across all simulated scenarios, to test the possible effects of Flood-MAR on additional leaching risk. For the model, planting occurred on April 15 every year, fertilization was applied as urea ammonium nitrate (UAN) in four splits with 50 kg total N applied in a preplant on April 15, and the remainder was divided evenly as 66.7 kg N applications on June 1, July 1, and July 21. UAN applications were entered into RZWQM2 input files as appropriate proportions of NH₄, NO₃⁻, and urea. Irrigation was managed using the automated option in RZWQM2, where irrigation is triggered when plant-available water is depleted to some allowable percentage threshold, called “allowable depletion,” and then the soil is refilled by irrigation back to field capacity. Allowable depletion was set at 60% during the first 20 days of crop growth, then reduced to 50% until harvest. This typically resulted in an irrigation frequency of several days during initial stand establishment and then every 5–14 days during the summer, depending on the soil's water holding capacity. Irrigation depths varied dynamically with rooting depth as the crop developed, reaching 5–8 cm per irrigation mid-summer. An irrigation application of 1.25 cm at planting was also made to stimulate germination at a uniform date. The simulated tillage depth was 15 cm with a tandem disk on April 15 before planting and then on September 15 and 16 after harvest with a bedder ridge operation on September 17. Harvest date varied

dynamically across years and scenarios and was defined as the removal of aboveground biomass with 10 cm stubble height and occurring at growth stage 0.95 (0–1) with 95% efficiency.

2.2 | Climate data

Five stations from the California Irrigation Management Information System (CIMIS) were chosen on a wet-to-dry precipitation gradient (569–159 mm year⁻¹) with at least a 37-year, largely intact record to provide climate data as input to RZWQM2 simulations (Table 1). Daily aggregated weather data were downloaded using the *cimir* package in R software (Koochafkan, 2021) and then subjected to in-house R software functions for visualizing anomalies and also automated quality control checks, error correction, and gap-filling procedures. Specifically, thresholds for possible minimum and maximum temperatures, relative humidity, insolation, wind, and precipitation were defined for each station based on official Fresno and Sacramento, CA, National Weather Service records and then corrected when necessary, using calculated daily means for the station. In some cases, missing precipitation or likely erroneous precipitation data were gap-filled using daily PRISM (parameter-elevation regressions on independent slopes model) data downloaded for the 4 km cell within which each CIMIS station was located. Additionally, some anomalous growing season precipitation values at the Five Points and Parlier stations, attributed to malfunctioning irrigation systems used to maintain a reference green grass surface surrounding the CIMIS weather stations, were corrected with PRISM data. For the Shafter station, erroneous precipitation data from 2004 to 2005 and multi-month missing data in 2012–2013 were provided by the nearby Delano CIMIS station 182. Finally, all temperature data were additionally corrected using daily means if a daily maximum or minimum temperature exceeded 3.5 standard deviations from its respective daily mean.

2.3 | Soil data

Note that 33 soil profile datasets from SSURGO were used to parameterize the 990 RZWQM2 scenarios (one BAU and five Flood-MAR scenarios across 165 soil × climate conditions). The soil data objective was to represent widespread soils without restrictive horizons, spanning a broad textural gradient, and to generally test whether differences in inherent soil physical properties affected NO₃⁻ leaching response to Flood-MAR. The SSURGO sampling strategy was based initially on a soil classification study that defined seven generalized soil regions in California. Of these seven, three typically well-drained soil regions varying in texture were used as sampling targets, including coarse soils with no

restrictions, loamy soils with no restrictions, and a fine-textured endmember, shrink-swell soils (Devine et al., 2021). The most widespread soil map unit components within each class were selected from the SSURGO with at least 80% assignment to one of these three soil regions used to create a sampling subset. Then, one representative soil profile dataset was selected randomly for each of the 11 most widespread soil components in each of the three soil regions. Only soil profiles with standard horizon nomenclature were sampled (A, B, C, etc.), since some SSURGO soil components from vintage survey areas have H1, H2, H3, and so forth nomenclature that indicates that soil data were entered without regard to a soil's typical horizonation, due to database constraints at the time of data entry.

The soil regions above were also found to correspond very well to generalized USDA-NRCS taxonomic family particle-size definitions (Soil Survey Staff, 1999). These USDA-NRCS soil classification definitions were ultimately used for the purpose of aggregating the results of this study as follows: coarse soils included those in the “coarse-loamy” and “sandy” classes, loamy soils included those in the “fine-loamy” and “fine-silty” classes, and fine soils included those in the “fine” and “very fine” classes. However, the other four soil regions in the seven-region soil classification approach, including soils with restrictive horizons and salt-affected soils (Devine et al., 2021), would also fall under these USDA-NRCS taxonomic particle-size classes, highlighting that not all of the Central Valley soil diversity has been represented in this study.

2.4 | Definition of “additional NO₃⁻ leaching risk”

For each unique Flood-MAR scenario (33 soils × 5 climates × 5 Flood-MAR timing strategies = 825 Flood-MAR scenarios), an “additional NO₃⁻ leaching risk” was calculated by subtraction of the total nitrate leached from its respective BAU scenario that used the same soil and climate (BAU for 33 soils × 5 climate combinations = 165 BAU scenarios). This relative “additional NO₃⁻ leaching risk” term is used throughout Section 3 and acknowledges that the study is not attempting to predict any absolute amount of NO₃⁻ leaching for a given place or time. The purpose of the BAU scenario is to serve as a control to measure the extent to which Flood-MAR impacts NO₃⁻ leaching. In other words, the interpretation of this study's results is meant to ascertain the relative risks of practicing Flood-MAR across different soils and climates in the Central Valley. The simulated maize cropping system with winter fallow is meant to serve as an analogue for any cropping system in California with a substantial dormant or fallow season, receiving ample N fertilizer applications with the potential for residual NO₃⁻ accumulation.




FIGURE 1 Preliminary (end of simulated initial conditions [SIC]) and final residual soil nitrate (in the business-as-usual scenario, no Flood-MAR) by three generalized taxonomic family particle-size classes: coarse, loamy, and fine ($n = 11$ soils per class). Each plot shows the different climates modeled from wettest (Durham, 537 mm year^{-1}) to driest (Shafter, 143 mm year^{-1}). Preliminary is the end of a 37-year preliminary run defining SIC. Final is the end of a subsequent 37-year run, using the SIC end-of-run values for initialization. Note the difference in Y-axis scale in wet versus drier sites. Box plots display the median and interquartile range. Whiskers display the distribution of data extending 1.5 times the upper and lower quartiles. Open circles are outliers.

3 | RESULTS AND DISCUSSION

3.1 | Wetter climates prevent accumulation of residual NO_3^- in business-as-usual scenarios

Under BAU scenarios, the end-of-run residual soil NO_3^- largely stabilized across generalized taxonomic family particle-size classes in each climate modeled (Figure 1). This is inferred by the similarity in residual nitrate levels between the SIC (after 37 years of “spin-up”) and at the end of the 37-year BAU simulation (Figure 1). This suggests that, for each unique soil and climate combination, the SIC represent the appropriate conditions for the dynamic steady-state residual NO_3^- in the modeled agroecosystem, thus allowing for a robust test of the effect of Flood-MAR on additional NO_3^- leaching risk. Whole soil residual NO_3^- in wetter climates

(Durham and Davis) were typically $60\text{--}100 \text{ kg N ha}^{-1}$ in the BAU scenario (Figure 1). Thus, climates with typical precipitation amounts of $>400 \text{ mm year}^{-1}$ (Table 1) were sufficient to leach soils in this simulated, fertilized agroecosystem, suggesting that Flood-MAR practiced in wetter Central Valley climates is of relatively lower additional NO_3^- leaching risk (Figure 2). Modeled residual NO_3^- levels and the relationship with mean annual precipitation generally corresponded with levels found in the literature, although we were unable to find data for California. In Minnesota, as an example, residual NO_3^- -N under corn production ranged from around 100 kg N ha^{-1} at locations with relatively low precipitation (400 mm year) to 42 kg N ha^{-1} at locations with high precipitation (891 mm year) (Yadav, 1997). In addition, a 4-year trial in corn in eastern Colorado showed gradual buildup ($23 \text{ kg ha}^{-1} \text{ NO}_3^-$ -N) of residual nitrate in plots that were suboptimally




FIGURE 2 Effect on 37-year, cumulative nitrate leached of different Flood-MAR strategies, including 3-day and 7-day frequency events practiced either in January or March or 21-day frequency events spanning early January through late March. Additional nitrate leached is relative to each soil's business-as-usual run (no Flood-MAR). Each plot is grouped by three generalized taxonomic family particle-size classes: coarse, loamy, and fine ($n = 11$ soils per class), and shows the different climates modeled from wettest (Durham) to driest (Shafter). See Table 1 for descriptions of climates. Note the difference in Y-axis scale in wet versus drier sites. Box plots display the median and interquartile range. Whiskers display the distribution of data extending 1.5 times the upper and lower quartiles. Open circles are outliers.

irrigated (46 cm of applied water) compared to half as much nitrate buildup ($13 \text{ kg ha}^{-1} \text{ NO}_3^- \text{-N}$) in plots that were over irrigated (85 cm of applied water) (Ludwick et al., 1976).

3.2 | Accumulation of residual NO_3^- in dry climates drives additional NO_3^- leaching risk

Residual soil NO_3^- increased by an order of magnitude in BAU coarse and loamy soils, but not in fine soils from wettest to driest climates (Figure 1). Compared to coarse soils, loamy soils tended to accumulate even more residual soil NO_3^- and displayed more variance from wet to dry. In the second driest climate (Five Points), BAU residual NO_3^- exceeded $500 \text{ kg NO}_3^- \text{-N ha}^{-1}$ in 5 of 11 coarse soils (interquartile range: $277\text{--}686 \text{ kg NO}_3^- \text{-N ha}^{-1}$), 8 of 11 loamy soils (interquar-

tile range: $498\text{--}1493 \text{ kg NO}_3^- \text{-N ha}^{-1}$), and 2 of 11 fine soils (interquartile range: $77\text{--}301 \text{ kg NO}_3^- \text{-N ha}^{-1}$). In the driest climate (Shafter), BAU residual NO_3^- exceeded $500 \text{ kg NO}_3^- \text{-N ha}^{-1}$ in 7 of 11 coarse soils (interquartile range: $370\text{--}882 \text{ kg NO}_3^- \text{-N ha}^{-1}$), 10 of 11 loamy soils (interquartile range: $586\text{--}1860 \text{ kg NO}_3^- \text{-N ha}^{-1}$), and 2 of 11 fine soils (interquartile range: $137\text{--}342 \text{ kg NO}_3^- \text{-N ha}^{-1}$). In the driest climate, extreme values of residual NO_3^- accumulated in 2 of 11 coarse soils ($1693\text{--}2249 \text{ kg NO}_3^- \text{-N ha}^{-1}$) and 4 of 11 loamy soils ($1831\text{--}3145 \text{ kg NO}_3^- \text{-N ha}^{-1}$), but the most extreme fine soil accumulated far less ($887 \text{ kg NO}_3^- \text{-N ha}^{-1}$).

Field studies in the region have demonstrated similar findings to this modeling exercise where soil profile nitrate was shown to accumulate during dry years and leach into the vadose zone during wet years (Liang et al., 1991; Rajj-Hoffman et al., 2024). Soil coring studies from the southern

portions of the San Joaquin Valley, California, which correspond with the driest sites in this study, show residual NO_3^- masses similar to but slightly lower ($180\text{--}269 \text{ kg ha}^{-1}$) than the range in values derived from these RZWQM2 simulations of fertilized maize (Bachand et al., 2014; Harter et al., 2005; Waterhouse et al., 2021). These studies sampled sites growing almonds, nectarines, tomatoes, grapes, and wheat, and did not span the gradient in texture modeled here.

In drier climates, loamy soils tended to present the greatest possibility of risk of additional NO_3^- leaching with Flood-MAR (Figure 2). In the driest climate (Shafter), 4 of 11 loamy soils leached $>3000 \text{ kg additional NO}_3^- \text{-N ha}^{-1}$ over 37 years under the 21-day frequency Flood-MAR scenario, while the median flux was $1270 \text{ kg additional NO}_3^- \text{-N ha}^{-1}$ (interquartile range: $665\text{--}3347 \text{ kg additional NO}_3^- \text{-N leached ha}^{-1}$). Although coarse- and fine-textured soil groups only leached marginally less NO_3^- (median) compared to loamy soils during Flood-MAR treatments (Figure 2), the interquartile ranges of coarse and fine textural groups were much narrower: $695\text{--}1007$ and $622\text{--}1214 \text{ kg additional NO}_3^- \text{-N leached ha}^{-1}$, respectively. In the second driest climate (Five Points), 3 of 11 loamy soils leached $>3000 \text{ kg additional NO}_3^- \text{-N ha}^{-1}$, with median fluxes of $990 \text{ kg additional NO}_3^- \text{-N ha}^{-1}$ (interquartile range: $609\text{--}2360 \text{ kg additional NO}_3^- \text{-N leached ha}^{-1}$). Similar to the driest climate, interquartile ranges of coarse and fine soil groups were much narrower: $499\text{--}764$ and $532\text{--}1103 \text{ kg additional NO}_3^- \text{-N leached ha}^{-1}$, respectively.

Although there are several facets of the N-cycle, which can complicate this risk assessment (Figures 3 and 4; Figures S1–S4), the most direct mechanistic explanation for additional NO_3^- leaching risk from loamy soils in the drier climates is due to their moderate level of microporosity and capacity to accumulate NO_3^- due to high water storage capacity. Loamy soils require more percolating water to leach effectively compared to coarse soils, explaining their conduciveness to residual NO_3^- accumulation in drier settings relative to coarse soils (Figure 1). These model findings are supported by deep core analysis from California field crops where a significant negative relationship was found between clay content in the soil profile (particle size control section) and NO_3^- concentration in the vadose zone ($1.8\text{--}8 \text{ m}$; Lund et al., 1974). The study found low residual NO_3^- concentrations ($<5 \mu\text{g g}^{-1}$) in the upper 1.8 m of sandy soils underlain by higher concentrations ($20 \mu\text{g g}^{-1}$) in the deep vadose zone. In contrast, comparatively high amounts of NO_3^- were found in the upper 1.8 m of loamy soils (peaking at $15 \mu\text{g g}^{-1}$) with lower concentrations ($5 \mu\text{g g}^{-1}$) in the underlying deep vadose zone. Onsoy et al. (2005), in a 15 m deep, highly heterogeneous vadose zone, reported the lowest $\text{NO}_3^- \text{-N}$ concentrations in coarse-textured sediment facies relative to finer textured sediments. Similarly, lab-based core leaching experi-

ments have shown finer textured soils retained more $\text{NO}_3^- \text{-N}$ compared to sandy soils (Gaines & Gaines, 1994).

While fine-textured soils also have high water storage capacity, residual N buildup was less (Figure 1) due to substantially higher denitrification (Figures 3 and 4). Fine-textured soils have the potential to experience higher denitrification rates and/or longer episodes of denitrification due to longer durations of elevated water-filled pore space (Groffman & Tiedje, 1988; Schindlbacher et al., 2004).

The RZWQM2 irrigation module uses an idealized irrigation schedule assuming perfect foresight by the farmer and uniform irrigation conditions: irrigation was triggered to apply water when a set threshold of plant-available water was depleted. Upon triggering, RZWQM2 applied a depth of water equal to refilling the soil's real-time, dynamic rooting depth to field capacity. This highly efficient irrigation decision approach had the effect of minimizing deep percolation, thus overestimating the accumulation of residual nitrate. To some extent, the higher NO_3^- mass leaching in the drier modeled climates would have been mitigated if this ideal BAU irrigation scheme was less efficient or if winter water applications were used to mitigate salt buildup, and thus lessen the impact of Flood-MAR on NO_3^- leaching concentrations.

Under real, practical conditions, conventional irrigation management approaches recommend applying more water than necessary to avoid creating soil moisture stress. Furthermore, irrigation systems do not apply water uniformly, and infiltration rates are inherently variable (Letey, 1985). For example, if 5 cm of applied water is needed to offset evapotranspiration, then a system with 80% irrigation distribution uniformity would need a 6.25-cm water application to prevent soil moisture stress. The consequence of this irrigation strategy is that most portions of a field will actually receive a depth of water greater than the amount necessary to refill the depleted soil moisture, resulting in deep percolation and NO_3^- leaching below the root zone as a consequence. A similar outcome is practiced in areas prone to salinity, where leaching fractions are used to manage the salinity of root zones (Rhoades et al., 1973), now often postharvest to avoid reducing nitrogen use efficiency. Applying more water to address salinity or irrigation distribution uniformity could complicate the interpretation of this study's results. However, while such real-world practices may mean more deep percolation than this study suggests, it would also mean the Flood-MAR risk for additional NO_3^- leaching in this study is a conservatively high "book-end" estimate. Our results present a relatively conservative risk assessment, and Flood-MAR would more clearly be a safe groundwater-quality mitigation tool in drier climates as well. Conversely, widespread use of subsurface drip irrigation throughout California has been shown to limit root zone leaching and/or minimize leaching to concentrated areas surrounding the drip line (Hanson et al., 2008), leading to conditions more closely reflecting the simulated BAU here.




FIGURE 3 (a–h) RZWQM2 simulated effects, comparing agricultural management of floodwaters (Flood-MAR) with business-as-usual (Control) on annualized N mass balance, hydrology, and water quality: (a) nitrate leached mass, (b) net mineralization, (c) harvested crop biomass N, (d) soil organic matter change affecting N balance, (e) volatilized N, (f) runoff, (g) deep percolation, and (h) NO_3^- in deep percolation. Each plot is grouped by generalized taxonomic family particle-size classes: coarse, loamy, and fine ($n = 11$ per class). The crop system is $250 \text{ kg N ha}^{-1} \text{ year}^{-1}$ fertilized and irrigated maize in Parlier, CA (median precipitation = 278 mm year^{-1}). Box plots display the median and interquartile range. Whiskers display the distribution of data extending 1.5 times the upper and lower quartiles. Open circles are outliers.

3.3 | Additional NO_3^- leaching risk in drier climates is mitigated by denitrification in finer textured soils and relative ease of leaching in coarse soils

The residual NO_3^- interquartile ranges of coarse and loamy texture groups tended to increasingly overlap from dry to driest (Figure 1). This convergence partly reflected the denitrification capacity of loamy soils under high residual NO_3^- conditions (Figure 4), which contributed to mitigate additional NO_3^- leaching risk of loamy soils in drier climates (Figure 2). However, loamy soils maintained an environment less favorable for denitrification compared to fine-textured soils (D’Haene et al., 2003), as simulated by RZWQM2 (Figure 4).

In fine-textured soils, denitrification limited the accumulation of residual soil NO_3^- to a greater extent, especially when dry climates favored buildup of NO_3^- due to less deep percolation (Figures 1, 3g, and 4; Figures S3g and S4g). Coarse soils displayed only negligible denitrification capacity, even under the driest climate (Figure 4). In fact, their capacity to accumulate some additional residual NO_3^- in the driest climate was reflected in the BAU run compared to the SIC run that assumed BAU conditions (Figure 1). In addition to deni-

trification, the greater microporosity of fine soils physically limited BAU NO_3^- leaching, especially in drier climates, making the notably lower total NO_3^- leached in Flood-MAR scenarios from fine soils more impactful on additional NO_3^- leaching risk due to Flood-MAR (Figures 3a and 5–7; Figures S1a–S4a). This is in spite of the fact that deep percolation depths from fine soils were not markedly different compared to coarse and loamy soils in drier climates (Figure 3g; Figures S3g–S4g). The retention of NO_3^- in soil under BAU, drier climate scenarios (Figures 5 and 7) helps explain the similar relative responsiveness of coarse and fine soils to Flood-MAR in terms of additional NO_3^- leached (Figure 2). In the BAU scenario under the driest climate, coarse soils typically leached $1779 \text{ kg NO}_3^- \text{-N ha}^{-1}$ (interquartile range: $1305\text{--}1917 \text{ kg NO}_3^- \text{-N ha}^{-1}$), compared to $117 \text{ kg NO}_3^- \text{-N ha}^{-1}$ (interquartile range: $37\text{--}327 \text{ kg NO}_3^- \text{-N ha}^{-1}$) from fine soils (Figures 5 and 7). Thus, this directly offset the Flood-MAR effect on leaching in coarse soils but exacerbated the NO_3^- leaching risk from fine soils, in spite of their high levels of denitrification and lower residual NO_3^- . Denitrification losses can be very high in fields with clay texture (Pratt et al., 1972), and the relationship between texture and denitrification has been documented where denitrification potential has been shown to be highest in fine-textured soils, moderate in loamy




FIGURE 4 Effect on 37-year, cumulative denitrification of business-as-usual (no Flood-MAR) and various Flood-MAR strategies, including 3-day and 7-day frequency events practiced either in January or March or 21-d frequency events spanning early January through late March. Each plot is grouped by three generalized taxonomic family particle-size classes: coarse, loamy, and fine ($n = 11$ soils per class), and shows the different climates modeled from wettest (Durham) to driest (Shafter). See Table 1 for descriptions of climates. Box plots display the median and interquartile range. Whiskers display the distribution of data extending 1.5 times the upper and lower quartiles. Open circles are outliers.

soils, and low in coarse soils when water-filled pore space exceeds 60% (D'Haene et al., 2003). A study of deep cores across a texture gradient in California found residual nitrate concentrations decreased with increasing clay content. The study suggests that while clay-rich soils have the potential to build up in N via incomplete leaching, this N pool is removed by denitrification (Lund et al., 1974). Greater losses of NO_3^- via denitrification were simulated in fine (silt loam)-textured compared to coarse (sandy loam)-textured vadose zones during Flood-MAR using a reactive transport model (Waterhouse et al., 2021).

3.4 | NO_3^- leaching risk “tipping point”

Comparing BAU deep percolation rates versus additional NO_3^- risk highlights a clearly nonlinear relationship to

climate and, specifically, precipitation (Figure 8). The inflection or “tipping point” of this relationship is perhaps best exemplified by the Parlier climate (median annual precipitation: 278 mm year⁻¹), where coarse soils showed less additional NO_3^- leaching risk compared to loamy and fine soils (Figure 2). The Parlier climate had ample precipitation to effectively leach most coarse soils in the initial BAU scenario, preventing accumulation of residual NO_3^- , but not enough for most loamy soils (Figure 1). Accordingly, loamy soils leached more additional NO_3^- during Flood-MAR compared to coarse soils in the Parlier climate (Figure 2). Despite substantial denitrification limiting residual NO_3^- accumulation in fine soils in the Parlier climate (Figures 1 and 4), the typical fine-textured soil leached slightly more additional NO_3^- (460 additional kg NO_3^- -N ha⁻¹) compared to loamy soils (395 additional kg NO_3^- -N ha⁻¹) (Figure 2). However, loamy soils arguably still had the greater additional NO_3^- leaching risk in




FIGURE 5 Typical soil profile NO_3^- (top row) and total NO_3^- leached (bottom row) for a *coarse* (taxonomic family particle-size class) soil, comparing business-as-usual (no Flood-MAR) versus managed groundwater recharge (Flood-MAR(21d) = 60 cm year⁻¹ across four applications on 21-day intervals), practiced during the 10 wettest winters from Davis (left column), the second wettest climate modeled, through Shafter (right column), the driest climate modeled. Note the difference in the Y-axis scale in wet versus drier sites.




FIGURE 6 Typical soil profile NO_3^- (top row) and total NO_3^- leached (bottom row) for a *loamy* (taxonomic family particle-size class) soil, comparing business-as-usual (no Flood-MAR) versus managed groundwater recharge (Flood-MAR(21d) = 60 cm year⁻¹ across four applications on 21-day intervals), practiced during the 10 wettest winters from Davis (left column), the second wettest climate modeled, through Shafter (right column), the driest climate modeled. Note the difference in the Y-axis scale in wet versus drier sites.




FIGURE 7 Typical soil profile NO_3^- (top row) and total NO_3^- leached (bottom row) for a *fine* (taxonomic family particle-size class) soil, comparing business-as-usual (no Flood-MAR) versus managed groundwater recharge ((Flood-MAR(21d) = 60 cm year⁻¹ across four applications on 21-day intervals), practiced during the 10 wettest winters in Davis (left column), the second wettest climate modeled, through Shafter (right column), the driest climate modeled. Note the difference in the Y-axis scale in wet versus drier sites.




FIGURE 8 Relationship between total, 37-year deep percolation in the business-as-usual scenario (no Flood-MAR) and additional nitrate leached due to Flood-MAR practiced during the 10 wettest water years (October–September), applying 60 cm⁻¹ water year⁻¹ across four applications on 21-day intervals. The vertical red line indicates the total Flood-MAR water applied over the 37-year period. Data shown across all climates modeled ($n = 5$) and soils ($n = 33$), symbolized by climate. Additional nitrate leached is relative to each soil's business-as-usual simulation in a particular climate. See Table 1 for description of climates.

the Parlier climate due to their propensity for more extreme residual NO_3^- accumulation, despite the relatively high levels of BAU NO_3^- leaching (3 of 11 soils developed >1000 kg residual NO_3^- -N ha⁻¹ in Parlier; Figure 1). This was directly

linked to additional NO_3^- leaching with Flood-MAR that was skewed toward more extreme fluxes in loamy soils (Figure 2). Close examination of RZWQM2 breakthrough curves for these more extreme loamy soils indicated that they did not generate substantial deep percolation until at least the second 15-cm Flood-MAR event. This explains the propensity of loamy soil to accumulate residual NO_3^- in BAU scenarios in the just-dry-enough Parlier climate with not enough precipitation during even the wettest years to generate substantial deep percolation.

3.5 | Effect of Flood-MAR timing

Despite the marked increase in additional NO_3^- leaching with Flood-MAR in drier climates, no timing strategy was effective at appreciably mitigating this risk. However, there were consistent effects of late-season versus early-season Flood-MAR timing across different soils. Specifically, coarse and loamy soils experienced more additional NO_3^- leached during later season Flood-MAR scenarios (Figure 2, March 3-day and 7-day). Early-season Flood-MAR experienced reduced additional NO_3^- leached because mineralization rates are subdued at cooler temperatures (Miller & Geisseler, 2018). Moreover, Flood-MAR had the effect of reducing annual denitrification compared to BAU (Figure 4) by decreasing residual soil NO_3^- through leaching before the growing season when denitrification potential is greatest on account of warmer, more

biogeochemically active soils. In addition to the amounts of O_2 and available C in soil, residual NO_3^- concentration is a primary control of denitrification (Myrold & Tiedje, 1985).

3.6 | Flood-MAR following multiyear droughts presents some risk

Across the 37-year simulation with Flood-MAR during the 10 wettest winters, median Flood-MAR scenarios tracked BAU scenarios closely in wetter climates: the typical coarse soil leached just 43 and 45 kg additional NO_3^- -N ha^{-1} in Durham and Davis, respectively (Figure 5); the typical loamy soil leached just 54 and 63 kg additional NO_3^- -N ha^{-1} in Durham and Davis (Figure 6); and the typical fine soil leached 152 and 336 additional NO_3^- -N ha^{-1} in Durham and Davis, respectively (Figure 5).

However, in drier climates, residual NO_3^- accumulated during periodic droughts in the climate record (e.g., 2012–2016), which was then flushed as deep percolation during subsequent wet years (e.g., 2017) in the Flood-MAR scenarios on typical coarse and loamy soils (Figures 5 and 6). Thus, the timing of Flood-MAR following multiyear droughts is of special concern in drier climates (median precipitation < 400 mm) and especially on more difficult-to-leach soils where high levels of residual NO_3^- accumulation are possible. It is during these dry years that residual NO_3^- would be accumulating, but it is relatively rare in these locations that wet-year precipitation is sufficient to initiate substantial deep percolation capable of leaching soils effectively. If precipitation whiplash continues, strategies will be needed to deplete residual nitrate either via improved fertigation methods (higher nitrogen use efficiency), crop rotations, or use of cover crops (Delgado et al., 1999).

In California, nutrient management plans are used as tools to help growers reduce nitrogen export from irrigated agriculture into groundwater and surface waters of the State. Nutrient management plans are designed to help growers understand the nitrogen cycle and the N budget, especially the amount of N left in soil after harvest. N budgeting may partially mitigate concerns about residual NO_3^- buildup in soil, which drives the outcomes of this study. Improved N management in the future may reduce the risk of leaching when Flood-MAR is practiced. However, this study simulated ideal irrigation and N application rates, which is a conservative estimation. Greater adoption of N budgeting would only make this study more relevant to actual field practices.

3.7 | Simulated runoff behavior in wetter climates partially influences results

The greater microporosity of higher clay content soils also explained why fine soils had the highest levels of additional

NO_3^- leached with Flood-MAR in the wetter climates of Durham and Davis, though the risk was relatively low compared to drier climates (Figure 2). The precipitation regimes of wetter climates were not completely effective at leaching the fine soils. However, the leaching response of two fine outlier soils in wetter climates was explained by their simulated high runoff, reducing effective precipitation and deep percolation in BAU scenarios (Figures S1–S3). Irrigation as applied in RZWQM2 cannot induce runoff (all applied water is assumed to infiltrate), which was the mechanism by which water was applied in the Flood-MAR scenarios. This was accepted as a realistic assumption because farmers practicing Flood-MAR would be expected to control runoff and manage for maximizing infiltration. In fact, to represent a comparable BAU scenario by minimizing the effect of runoff in RZWQM2 in wet climates, daily precipitation data from CIMIS were assumed to occur in 12-h storms with variable precipitation intensities of 0.05 (min) – 5.0 (max) $cm\ h^{-1}$, creating low runoff conditions in BAU scenarios for most soils (Figure 4; Figures S1–S4). On the other hand, if an actual “business-as-usual” scenario involved a field with, for example, surface sealing and infiltration issues, resulting in appreciable runoff and reduced deep percolation, then a transition to Flood-MAR management aimed at improved infiltration and enhanced deep percolation could present additional NO_3^- leaching risk not represented in this study.

3.8 | Simulated soil organic matter accumulation partially influences additional NO_3^- leaching risk, especially in coarse soils

RZWQM2 simulations consistently projected greater SOM accumulation in the coarse soil scenarios and hence, greater soil organic N accumulation (Figure 3d; Figures S1–S4). The change in organic N was above 15 kg N ha^{-1} in coarse soils compared to less than 5 kg N ha^{-1} in loamy and fine soils. This phenomenon reduced the total N pool available for leaching from coarse soils across growing seasons. This also partly explained the higher additional NO_3^- leaching risk under loamy soils compared to coarse soils.

3.9 | Implications for groundwater and well water nitrate concentrations

Results shown here are relevant due to the potentially long-term implementation of various Flood-MAR regimes as climate resiliency and groundwater (supply) sustainability practices are expected to expand in the future (Marr et al., 2018). Potentially widespread implementation of these practices is being considered to maximize the retention of floodwaters in aquifers rather than their loss to the ocean

(Marr et al., 2018). Simulations here pertain to losses of NO_3^- from the most dynamic and reactive portion of the unsaturated zone, the upper 2 m of soil. Conservatively, assuming relatively low reactivity in the vadose zone below soil and furthermore assuming laterally widespread application of these simulated conditions, the NO_3^- concentration in leachate from soil would be similar when it becomes recharge at the water table. And, if that recharge has the same concentration across a relatively large lateral extent, it would be expected that shallower (predominantly domestic) wells also experience concentrations of that magnitude, albeit after some period of delay corresponding to the travel time from the root zone to the water table and from the water table to the well (typically a few to tens of years). In this section, we address two questions: what concentration of NO_3^- do the aforementioned additional losses of NO_3^- mass due to Flood-MAR represent? And, second, how is that concentration different from those under BAU conditions?

The difference in mass losses from soil ranges from a few tens to several thousand kg N ha^{-1} over the simulated period (Figures 5–7). The main difference in water percolation over that same period is approximately 600 cm across all scenarios (the amount of additional Flood-MAR applied). The drinking water limit of $10 \text{ mg NO}_3^- \text{-N L}^{-1}$ is equivalent to $1 \text{ kg N}/(\text{ha}\cdot\text{l cm})$ or 10 kg N ha^{-1} in 10 cm of water (10 kg N in $0.1 \text{ ha}\cdot\text{m}$ of water) or 600 kg N ha^{-1} in 600 cm of water. From basic mass balance considerations, the *additional* water percolating from Flood-MAR does not exceed the nitrate maximum contaminant level (MCL), where the additional N leaching over the simulation period does not exceed 600 kg N ha^{-1} (Figure 8: almost all wet climate scenarios in Durham and Davis, some scenarios in the intermediate climate in Parlier, and few scenarios in the dry climate). Under dry climate conditions, where median N losses are on the order of twice 600 kg N ha^{-1} , the *additional* water percolating from Flood-MAR may be on the order of twice the MCL concentration, in some cases five times the MCL concentration (where the additional leaching is $3000 \text{ kg N ha}^{-1}$). Clearly, under the drier climate conditions, without adding additional practices such as improved nitrogen use efficiency, crop rotation, cover crops, or higher Flood-MAR applications than 60 cm in the wettest winters, the percolation due to the additional Flood-MAR water cannot meet drinking water standards. In fact, modeled NO_3^- concentration in deep percolation waters receiving Flood-MAR treatment was reduced to equal the drinking water standard for most fine soil groups, while being slightly to moderately above the standard for loamy and coarse texture groups depending on location (Figure 3h; Figures S1–S4).

The NO_3^- concentrations of deep percolation water in the BAU scenarios were higher in dry regions (Figure 3h). Moreover, NO_3^- concentrations of deep percolation water in

the BAU scenarios can be estimated from the total mass of N leached under BAU conditions (no Flood-MAR; Figures 5–7) and the corresponding total BAU percolation (Figure 8). The total BAU NO_3^- leached was highest in coarse soils, ranging from $1000 \text{ kg N ha}^{-1}$ under dry climates to over $1500 \text{ kg N ha}^{-1}$ in wet climates (Figure 5). The corresponding percolation rates over the simulation period range from less than 20 to 250 cm in drier climates and from 300 to over 1200 cm in wet climates (Figure 8). Hence, under BAU conditions of drier climate, nitrate-N concentrations in (very slowly) percolating water exceed the MCL by nearly one to nearly two orders of magnitude ($1500/250$ – $1500/20 \text{ kg N}/(\text{ha}\cdot\text{cm water})$), while, under BAU conditions of wetter climates, nitrate-N concentrations in (rapidly) percolating water range from at the MCL to five times above the MCL ($1500/300$ – $1500/1200 \text{ kg N}/(\text{ha}\cdot\text{cm water})$) under the simulated conditions.

Hence, under all climate conditions, the NO_3^- concentration in the *additional* percolating water due to Flood-MAR is significantly lower than the NO_3^- concentration in percolating water from BAU conditions. Due to the mixing of these two waters in the root zone and further in the deeper unsaturated zone, shallow groundwater, and along the vertical well screen of an extraction well, the net long-term effect of Flood-MAR is a significant reduction of NO_3^- concentration in shallow groundwater and, hence, in domestic and shallow public water supply water under the dominant influence of agricultural return water recharge. Despite the fact that Flood-MAR leads to additional nitrate *mass* leaching, especially under dry climate conditions, the amount of Flood-MAR in the simulated scenarios is sufficient to dilute this additional nitrate mass to less than or slightly above the NO_3^- MCL depending on soil type and climate (Figure 3h). Pratt et al. (1972) arrived at a similar finding showing low $[\text{NO}_3^-]$ (below MCL) in deep leachate from orange orchards in the Central Valley due to the application of excess irrigation water to maintain low salt levels in soils.

The above considerations, while hypothetical, are highly relevant for consideration of the long-term impacts of Flood-MAR on well water NO_3^- concentrations but will be further modified due to the fact that stream and incidental canal recharge with typically negligible NO_3^- concentrations plays an important role particularly in the drier climate conditions of the southern Central Valley, leading to further dilution of irrigated lands return flow recharge NO_3^- in some wells. We note that, under the typical percolation and recharge rates of the drier climate scenarios ($50 \text{ cm}/37 \text{ years}$) when compared to the wet climate scenarios ($500 \text{ cm}/37 \text{ years}$; Figure 8), the source area of a well, at any given pumping rate, would be an order of magnitude larger in dry climate locations than in wet climate location, assuming the entire landscape was subject to the simulated scenario conditions.

4 | STUDY LIMITATIONS

This study evaluated the nitrate leaching risk of 33 different soil series with contrasting physical properties representative of commonly mapped soils across the region. The broad scale of the study domain, the magnitude of time evaluated, and the large number of Flood-MAR scenarios investigated made it impossible to calibrate and validate the model uniformly with field data. Thus, for the most part, model default conditions were used. Model calibration is site-specific, and this study lacked representative locations for this purpose. As a result, results reflect state factor controls (climate and soil properties that influence the water balance) on additional NO_3^- leaching risk through a theoretical representation of N cycling in response to Flood-MAR scenarios. Future studies could explore improvements in model simulations resulting from site-specific calibrations.

Similarly, the study used soil survey data to parameterize physical and hydraulic model conditions. RZWQM2 uses the Brooks and Corey parameters to characterize soil water retention. This approach has been widely tested in diverse settings and soil types (Ahuja et al., 2000; Ma et al., 2006). More recent studies, however, have shown that optimization procedures to calibrate hydraulic parameters in the model using Latin hypercube sampling and gradient-based optimization improved the model performance (Fang et al., 2010; Ma et al., 2011). The results of our approach could have higher uncertainty compared to in-depth parameterization procedures. Moreover, the depth of soils studied here is not equal, varying randomly between 150 and 218 cm, which may have influenced some of the variability seen among texture classes.

The study did not assume surface ponding during Flood-MAR scenarios, which could occur. Ponding has the potential to increase the rate of deep percolation when the soil is saturated. Ponding is likely to have minimal impact on denitrification rates because the process is occurring across a time frame spanning soil water states from saturation to field capacity when air-filled pore space is low (Bateman & Baggs, 2005; Mekala & Nambi, 2017). In addition, the fate of NO_3^- in soil during the dormant season is largely controlled by temperature and the amount of water applied, not the differences in flow rate as influenced by ponded versus non-ponded conditions (Ludwick et al., 1976; Miller & Geiseler, 2018; Or et al., 2007). Moreover, our simulations, which reflect different durations of saturation, show no difference in denitrification when comparing 3-, 7-, or 21-day Flood-MAR treatments. This indicates that denitrification during Flood-MAR periods is controlled by low temperature and the initial amount of NO_3^- in the soil rather than the duration of saturation. Moreover, ponding is likely to be discouraged during Flood-MAR due to its potential harm to crops and surface soil condition (O'Geen et al., 2015). We perceive and recommend

Flood-MAR to resemble water application strategies similar to surface irrigation procedures in order to limit the negative effects of standing water.

This study only modeled one of the over 200 crops grown in California. Various crops will inevitably have different N requirements, irrigation strategies, rooting depths, and N use efficiencies, which will influence residual nitrate buildup. Although the residual N accumulation will differ among crops, this study is generalizable to any crop that has a dormant period over the winter months and some degree of residual nitrate accumulation. Clearly, absolute amounts of residual N buildup will differ by fertilization rate and crop uptake, and, most importantly, irrigation efficiency, but the climate and soil are what drive the buildup. Corn as a test crop is least generalizable to crops that have low N demand such as wine grapes and alfalfa.

Despite these limitations, this study has value in presenting new testable research questions and management considerations such as the likelihood of denitrification during the dormant season, the potential buildup of residual nitrate in response to drought, the effects of irrigation management, the potential impact of climate whiplash on residual NO_3^- , and the relevance of the N-cycle process in the context of Flood-MAR. Findings are not intended to prescribe the ideal conditions for Flood-MAR; rather, they justify the importance of documenting residual nitrate concentrations when considering this practice.

5 | CONCLUSION

Coarse soils are considered of the greatest risk to NO_3^- leaching into groundwater when N-fertilizer applications exceed N-removal processes in an agroecosystem: export in crop biomass, accumulation in SOM, volatilization, and denitrification (Figure 4). However, this truism did not hold up to evaluations of the effect of agricultural management of floodwater (Flood-MAR) on additional NO_3^- leaching risk. Simply put, except in the driest climates, precipitation is sufficient in California's Central Valley to leach residual NO_3^- , such that the additional NO_3^- leaching risk presented by Flood-MAR is typically lower in coarse soils compared to loamy soils.

Overall, this research suggests that Flood-MAR timing strategies, specifically seasonality and intervals between Flood-MAR events, are generally negligible in their effects on NO_3^- leaching risk in comparison to the risk of accumulating residual nitrate in dry climates. Nevertheless, the results point to some broader timing concerns for using Flood-MAR as a general strategy to recharge depleted aquifers. Most importantly, growers are advised to manage nitrogen carefully in dry years and monitor residual NO_3^- following successive

dry years, especially in locations where natural deep percolation is already typically limited by low rainfall. Residual NO_3^- may accumulate most rapidly during droughts, especially when growers face limited water resources that may typically result in lower-than-expected growing season soil flushing due to irrigation and leaching practices. However, under all scenarios, Flood-MAR is likely to noticeably reduce NO_3^- concentration in groundwater relative to BAU over the long run.

AUTHOR CONTRIBUTIONS

Scott Devine: Conceptualization; data curation; formal analysis; investigation; methodology; validation; visualization; writing—original draft; writing—review and editing. **Helen Dahlke:** Conceptualization; funding acquisition; writing—review and editing. **Thomas Harter:** Writing—review and editing. **Isaya Kisekka:** Writing—review and editing. **Majdi Abou Najm:** Writing—review and editing. **Anthony T. O’Geen:** Conceptualization; funding acquisition; investigation; project administration; resources; supervision; writing—original draft; writing—review and editing.

ACKNOWLEDGMENTS

Authors gratefully acknowledge the support of the California Department of Food and Agriculture–Fertilizer Research and Education Program.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflicts of interest.

ORCID

Scott Devine  <https://orcid.org/0000-0002-2306-8281>

Helen Dahlke  <https://orcid.org/0000-0001-8757-6982>

Isaya Kisekka  <https://orcid.org/0000-0002-2460-7777>

Majdi Abou Najm  <https://orcid.org/0000-0002-8434-2520>

Anthony T. O’Geen  <https://orcid.org/0000-0003-1499-511X>

REFERENCES

- Ahuja, L. R., Rojas, K. W., Hanson, J. D., Shaffer, M. J., & Ma, L. (Eds.). (2000). *Root zone water quality model*. Water Resources Publications.
- Bachand, P. A. M., Roy, S. B., Choperena, J., Cameron, D., & Horwath, W. R. (2014). Implications of using on-farm flood flow capture to recharge groundwater and mitigate flood risks along the Kings River, CA. *Environmental Science & Technology*, 48, 13601–13609. <https://doi.org/10.1021/es501115c>
- Bateman, E. J., & Baggs, E. M. (2005). Contributions of nitrification and denitrification to N_2O emissions from soils at different water-filled pore space. *Biology and Fertility of Soils*, 41, 379–388. <https://doi.org/10.1007/s00374-005-0858-3>
- Cameira, M. R., Fernando, R. M., Ahuja, L. R., & Ma, L. (2007). Using RZWQM to simulate the fate of nitrogen in field soil–crop environment in the Mediterranean region. *Agricultural Water Management*, 90(1–2), 121–136. <https://doi.org/10.1016/j.agwat.2007.03.002>
- CDWR. (2018). *Flood-MAR: Using flood water for managed aquifer recharge to support sustainable water resources*. California Department of Water Resources. https://cawaterlibrary.net/wp-content/uploads/2018/07/DWR_FloodMAR-White-Paper_06_2018_updated.pdf
- CDWR. (2022). *Groundwater sustainability plans*. California Department of Water Resources. <https://water.ca.gov/Programs/Groundwater-Management/SGMA-Groundwater-Management/Groundwater-Sustainability-Plans>
- Delgado, J. A., Sparks, R. T., Follett, R. F., Sharkoff, J. L., & Rigenbach, R. R. (1999). Use of winter cover crops to conserve water and water quality in the San Luis Valley of south central Colorado. In R. Lal (Ed.), *Soil quality and soil erosion* (pp. 125–142). CRC Press.
- Devine, S. M., Steenwerth, K. L., & O’Geen, A. T. (2021). A regional soil classification framework to improve soil health diagnosis and management. *Soil Science Society of America Journal*, 85, 361–378. <https://doi.org/10.1002/saj2.20200>
- D’Haene, K., Moreels, E., DeNeve, S., Daguilar, B., Boeckx, P., Hofman, G., & Van Cleemput, O. (2003). Soil properties influencing the denitrification potential of Flemish agricultural soils. *Biology and Fertility of Soils*, 38, 358–366.
- Domagalski, J. L., Phillips, S. P., Bayless, E. R., Zamora, C., Kendall, C., Wildman, R. A., & Hering, J. G. (2008). Influences of the unsaturated, saturated, and riparian zones on the transport of nitrate near the Merced River, California, USA. *Hydrogeology Journal*, 16, 675–690. <https://doi.org/10.1007/s10040-007-0266-x>
- Fang, Q.-X., Green, T. R., Ma, L., Erskine, R. H., Malone, R. W., & Ahuja, L. R. (2010). Optimizing soil hydraulic parameters in RZWQM2 using automated calibration methods. *Soil Science Society of America Journal*, 74, 1897–1913. <https://doi.org/10.2136/sssaj2009.0380>
- FAO. (2021). *World food and agriculture—Statistical yearbook 2021*. Food and Agriculture Organization of the United Nations. <https://doi.org/10.4060/cb4477en>
- Gaines, T. P., & Gaines, S. T. (1994). Soil texture effect on nitrate leaching in soil percolates. *Communications in Soil Science and Plant Analysis*, 25, 2561–2570. <https://doi.org/10.1080/00103629409369207>
- Geisseler, D., Lazicki, P. A., Pettygrove, G. S., Ludwig, B., Bachand, P. A. M., & Horwath, W. R. (2012). Nitrogen dynamics in irrigated forage systems fertilized with liquid dairy manure. *Agronomy Journal*, 104(4), 897–907. <https://doi.org/10.2134/agronj2011.0362>
- Green, W. H., & Ampt, G. A. (1911). Studies on soil physics: The flow of air and water through soils. *Journal of Agricultural Science*, 4(1), 1–24.
- Groffman, P. M., & Tiedje, J. M. (1988). Denitrification hysteresis during wetting and drying cycles in soil. *Soil Science Society of America Journal*, 52, 1626–1629. <https://doi.org/10.2136/sssaj1988.03615995005200060022x>
- Hanson, B., Hopmans, J. W., & Šimůnek, J. (2008). Leaching with subsurface drip irrigation under saline, shallow groundwater conditions. *Vadose Zone Journal*, 7, 810–818. <https://doi.org/10.2136/vzj2007.0053>
- Harter, T., Onsoy, Y., Heeren, K., Denton, M., Weissmann, G., Hopmans, J., & Horwath, W. (2005). Deep vadose zone hydrology demonstrates fate of nitrate in eastern San Joaquin Valley. *California Agriculture*, 59, 124–132. <https://doi.org/10.3733/ca.v059n02p124>
- Herbert, C., & Döll, P. (2019). Global assessment of current and future groundwater stress with a focus on transboundary aquifers.

- Water Resources Research*, 55, 4760–4784. <https://doi.org/10.1029/2018WR023321>
- Hoogenboom, G., Porter, C. H., Boote, K. J., Shelia, V., Wilkens, P. W., Singh, U., White, J. W., Asseng, S., Lizaso, J. I., Moreno, L. P., Pavan, W., Ogoshi, R., Hunt, L. A., Tsuji, G. Y., & Jones, J. W. (2019). The DSSAT crop modeling ecosystem. In K. J. Boote (Ed.), *Advances in crop modeling for a sustainable agriculture* (pp. 173–216). Burleigh Dodds Science Publishing. <https://dx.doi.org/10.19103/AS.2019.0061.10>
- Kisekka, I., Schlegel, A., Ma, L., Gowda, P. H., & Prasad, P. V. V. (2017). Optimizing preplant irrigation for maize under limited water in the high plains. *Agricultural Water Management*, 187, 154–163. <https://doi.org/10.1016/j.agwat.2017.03.023>
- Kocis, T. N., & Dahlke, H. E. (2017). Availability of high-magnitude streamflow for groundwater banking in the Central Valley, California. *Environmental Research Letters*, 12(8), Article 084009. <https://doi.org/10.1088/1748-9326/aa7b1b>
- Koohafkan, M. (2021). cimr: Interface to the CIMIS Web API (R Package Version 0.4-1) [Computer software]. CRAN. <https://CRAN.R-project.org/package=cimr>
- Letej, J. (1985). Irrigation uniformity as related to optimum crop production—Additional research is needed. *Irrigation Science*, 6, 253–263. <https://link.springer.com/article/10.1007/BF00262470>
- Levintal, E., Kniffin, M. L., Ganot, Y., Marwaha, N., Murphy, N. P., & Dahlke, H. E. (2022). Agricultural managed aquifer recharge (Ag-MAR)—A method for sustainable groundwater management: A review. *Critical Reviews in Environmental Science and Technology*, 53, 291–314. <https://doi.org/10.1080/10643389.2022.2050160>
- Liang, B. C., Remillard, M., & MacKenzie, A. F. (1991). Influence of fertilizer, irrigation, and non-growing season precipitation on soil nitrate-nitrogen under corn. *Journal of Environmental Quality*, 20, 123–128. <https://doi.org/10.2134/jeq1991.00472425002000010019x>
- Liu, P. W., Famiglietti, J. S., Purdy, A. J., Adams, K. H., McEvoy, A. L., Reager, J. T., Bindlish, R., Wiese, D. N., David, C. H., & Rodell, M. (2022). Groundwater depletion in California's Central Valley accelerates during megadrought. *Nature Communications*, 13, Article 7825. <https://doi.org/10.1038/s41467-022-35582-x>
- Ludwick, A. E., Reuss, J. O., & Langin, E. J. (1976). Soil nitrates following four years continuous corn and as surveyed in irrigated farm fields of central and eastern Colorado. *Journal of Environmental Quality*, 5(1), 82–86.
- Lund, L. J., Adriano, D. C., & Pratt, P. F. (1974). Nitrate concentrations in deep soil cores as related to soil profile characteristics. *Journal of Environmental Quality*, 3, 78–82. <https://doi.org/10.2134/jeq1974.00472425000300010021x>
- Ma, L., Ahuja, L. R., Saseendran, S. A., Malone, R. W., Green, T. R., Nolan, B. T., Bartling, P. N. S., Flerchinger, G. N., Boote, K. J., & Hoogenboom, G. (2011). A protocol for parameterization and calibration of RZWQM2 in field research. In L. R. Ahuja & L. Ma (Eds.), *Methods of introducing system models into agricultural research* (pp. 1–64). ASA, CSSA, SSSA. <https://doi.org/10.2134/advgricsystemmodel2.c1>
- Ma, L., Hoogenboom, G., Ahuja, L. R., Ascough, J. C., & Saseendran, S. A. (2006). Evaluation of the RZWQM-CERES-Maize hybrid model for maize production. *Agricultural Systems*, 87(3), 274–295. <https://doi.org/10.1016/j.agry.2005.02.001>
- Ma, L., Shaffer, M. J., Ahuja, L. R., Rojas, K. W., Xu, C., Boyd, J. K., & Waskom, R. (1998). Manure management in an irrigated silage corn field: Experiment and modeling. *Soil Science Society of America Journal*, 62, 1006–1017. <https://doi.org/10.2136/sssaj1998.03615995006200040023x>
- Marr, J., Arrate, D., Maendly, R., Dhillon, D., & Stygar, S. (2018). *FLOOD-MAR: Using flood water for managed aquifer recharge to support sustainable water resources* [White paper]. California Department of Water Resources. <https://water.ca.gov/Programs/All-Programs/Flood-MAR>
- Mekala, C., & Nambi, I. M. (2017). Understanding the hydrologic control of N cycle: Effect of water filled pore space on heterotrophic nitrification, denitrification and dissimilatory nitrate reduction to ammonium mechanisms in unsaturated soils. *Journal of Contaminant Hydrology*, 202, 11–22. <https://doi.org/10.1016/j.jconhyd.2017.04.005>
- Miller, K. S., & Geisseler, D. (2018). Temperature sensitivity of nitrogen mineralization in agricultural soils. *Biology and Fertility of Soils*, 54, 853–860. <https://doi.org/10.1007/s00374-018-1309-2>
- Murphy, N. P., Waterhouse, H., & Dahlke, H. E. (2021). Influence of agricultural managed aquifer recharge on nitrate transport: The role of soil texture and flooding frequency. *Vadose Zone Journal*, 20(5), 1–16. <https://doi.org/10.1002/vzj2.20150>
- Myrold, D. D., & Tiedje, J. M. (1985). Diffusional constraints on denitrification in soil. *Soil Science Society of America Journal*, 49, 651–657. <https://doi.org/10.2136/sssaj1985.03615995004900030025x>
- Nelson, R. L. (2012). Assessing local planning to control groundwater depletion: California as a microcosm of global issues. *Water Resources Research*, 48, Article W01502. <https://doi.org/10.1029/2011WR010927>
- Nimah, N. M., & Hanks, R. J. (1973). Model for estimating soil water, plant, and atmospheric interrelations: I. Description and sensitivity. *Soil Science Society of America Journal*, 37, 522–527. <https://doi.org/10.2136/sssaj1973.03615995003700040018x>
- O'Geen, A. T., Saal, M. B., Dahlke, H., Doll, D., Elkins, R., Fulton, A., Fogg, G., Harter, T., Hopmans, J. W., Ingels, C., Niederholzer, F., Sandovol, S., Verdegaaal, P., & Walkinshaw, M. (2015). Soil suitability index identifies potential areas for groundwater banking on agricultural lands. *California Agriculture*, 69, 75–84.
- Onsoy, Y. S., Harter, T., Ginn, T. R., & Horwath, W. R. (2005). Spatial variability and transport of nitrate in a deep alluvial vadose zone. *Vadose Zone Journal*, 4, 41–54. <https://doi.org/10.2136/vzj2005.0041a>
- Or, D., Smets, B. F., Wraith, J. M., Dechesne, A., & Friedman, S. P. (2007). Physical constraints affecting bacterial habitats and activity in unsaturated porous media—A review. *Advances in Water Resources*, 30, 1505–1527. <https://doi.org/10.1016/j.advwatres.2006.05.025>
- Pratt, P. F., Jones, W. W., & Hunsaker, V. E. (1972). Nitrate in deep soil profiles in relation to fertilizer rates and leaching volume. *Journal of Environmental Quality*, 1, 97–102. <https://doi.org/10.2134/jeq1972.00472425000100010024x>
- Qin, Y., Abatzoglou, J. T., Siebert, S., Huning, L. S., AghaKouchak, A., Mankin, J. S., Hong, C., Tong, D., Davis, S. J., & Mueller, N. D. (2020). Agricultural risks from changing snowmelt. *Nature Climate Change*, 10, 459–465. <https://doi.org/10.1038/s41558-020-0746-8>
- Raij-Hoffman, I., Dahan, O., Dahlke, H. E., Harter, T., & Kisekka, I. (2024). Assessing nitrate leaching during drought and extreme precipitation: Exploring deep vadose-zone monitoring, groundwater observations, and field mass balance. *Water Resources Research*, 60, Article e2024WR037973. <https://doi.org/10.1029/2024WR037973>

- Rawls, W. J., Brakensiek, D. L., & Saxton, K. E. (1982). Estimation of soil-water properties. *Transactions of the ASAE*, 25, 1316–1320. <https://doi.org/10.13031/2013.33720>
- Rhoades, J. D., Ingvalson, R. D., Tucker, J. M., & Clark, M. (1973). Salts in irrigation drainage waters: I. Effects of irrigation water composition leaching fraction, and time of year on the salt compositions of irrigation drainage waters. *Soil Science Society of America Journal*, 37, 770–774. <https://doi.org/10.2136/sssaj1973.03615995003700050038x>
- Richards, L. A. (1931). Capillary conduction of liquids through porous mediums. *Journal of Applied Physics*, 1, 318–333. <https://doi.org/10.1063/1.1745010>
- Schindlbacher, A., Zechmeister-Boltenstern, S., & Butterbach-Bahl, K. (2004). Effects of soil moisture and temperature on NO, NO₂, and N₂O emissions from European forest soils. *Journal of Geophysical Research*, 109, Article D17. <https://doi.org/10.1029/2004JD004590>
- Siebert, S., Burke, J., Faures, J. M., Frenken, K., Hoogeveen, J., Döll, P., & Portmann, F. T. (2010). Groundwater use for irrigation—A global inventory. *Hydrology and Earth System Sciences*, 14, 1863–1880. <https://doi.org/10.5194/hess-14-1863-2010>
- Soil Survey Staff. (1999). *Soil taxonomy: A basic system of soil classification for making and interpreting soil surveys* (Agricultural Handbook No. 436) (2nd ed.). USDA-NRCS.
- Swain, D. L., Langenbrunner, B., Neelin, J. D., & Hall, A. (2018). Increasing precipitation volatility in twenty-first-century California. *Nature Climatic Change*, 8, 427–33. <https://doi.org/10.1038/s41558-018-0140-y>
- Waterhouse, H., Arora, B., Spycher, N. F., Nico, P. S., Ulrich, C., Dahlke, H. E., & Horwath, W. R. (2021). Influence of agricultural managed aquifer recharge (AgMAR) and stratigraphic heterogeneities on nitrate reduction in the deep subsurface. *Water Resources Research*, 57, Article e2020WR029148. <https://doi.org/10.1029/2020WR029148>
- Waterhouse, H., Bachand, S., Mountjoy, D., Choperena, J., Bachand, P., Dahlke, H., & Horwath, W. (2020). Agricultural managed aquifer recharge—Water quality factors to consider. *California Agriculture*, 74(3), 144–154. <https://doi.org/10.3733/ca.2020a0020>
- Yadav, S. N. (1997). Formulation and estimation of nitrate-nitrogen leaching from corn cultivation. *Journal of Environmental Quality*, 26, 808–814. <https://doi.org/10.2134/jeq1997.00472425002600030031x>

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Devine, S., Dahlke, H., Harter, T., Kisekka, I., Najm, M. A., & O'Geen, A. T. (2025). Modeling nitrate leaching risk from flood managed aquifer recharge in California's agricultural lands. *Vadose Zone Journal*, 24, e70058. <https://doi.org/10.1002/vzj2.70058>